

On the Cubical Homology Groups of Free Partially Commutative Monoids

Ahmet A. Husainov

Abstract

We study a Leech homology of a locally bounded free partially commutative monoid $M(E, I)$. Given a contravariant natural system of abelian groups F on $M(E, I)$ we build a precubical set $T(E, I)$ with a homological system of abelian groups $\overline{F}$ and prove that the Leech homology groups $H_n(M(E, I), F)$ are isomorphic to the cubical homology groups $H_n(T(E, I), \overline{F})$, $n \geq 0$. As a consequence we have confirmed a conjecture that if the free partially commutative monoid does not contain $> n$ mutually commuting generators, then its the homological dimension $\leq n$. We have built the complexes of finite length for a computation of the Leech homology of such monoids and the Hochschild homology of their monoid rings. The results are applied to the homology of asynchronous transition systems. We give the positive answer to a question that the homological dimension of the asynchronous system does not greater than the maximal number of its mutually independent events. We have built the complex for computing the integral homology groups of an asynchronous transition system by the Smith normal form of integer matrices.

2000 Mathematics Subject Classification 16E40, 18B20, 18G10, 18G20, 18G35, 18G40, 20M50, 55U99, 68Q85

Keywords: cubical homology, precubical set, homology of small categories, Baues-Wirsching cohomology, Leech cohomology, Hochschild homology, asynchronous transition system.

Contents

1 Homology of categories and precubical sets

5

1.1	Precubical sets	5
1.2	Smith normal form and calculating the homology groups . . .	7
1.3	Homology of small categories	9
1.4	Homology of precubical sets	16
1.5	The category of factorizations	23
1.6	Acyclic partially ordered sets	25
2	Homology groups of free partially commutative monoids	26
2.1	Homology of a commutative monoid	26
2.2	Leech dimension of a free partially commutative monoid . . .	30
2.3	Leech homology groups of free partially commutative monoids	36
2.4	Hochschild homology	38
2.5	Homology with coefficients in a right module	40
3	Homology of right $M(E, I)$-sets with coefficients in diagrams	42
3.1	Homology groups of right $M(E, I)$ -sets	42
3.2	Homology of asynchronous transition systems	46

Introduction

This work contains three sections. In the first section we study homology groups of precubical sets with coefficients in homological systems of abelian groups. It is known [5, Application 2, Prop. 4.2] that the homology groups $H_n(X, F)$ of any simplicial set X with coefficients in a homological system of abelian groups F are isomorphic to the values $\varinjlim_n^{(\Delta/X)^{op}} F$ of the left satellites of the colimit where Δ/X is the category of singular simplexes of X . Similar assertion is true for the homological systems on semisimplicial sets (see [18, Prop. 1.4] for the dual assertion). It does not true in case of cubical sets (see Example 1.12). Nevertheless the main result of the first section (Theorem 1.13) shows that it is true for homological systems on the precubical sets. This allows us to use the homology theory of small categories. Using Oberst's Theorem [26, Theorem 2.3], we find the criterion under which the homomorphisms of homology groups of precubical sets corresponding to the precubical morphism are isomorphisms. We will build spectral sequences of a locally ordered covering (Cor. 1.16) and of a morphism of precubical sets (Cor. 1.17).

Second section is devoted to the homological dimension of monoids satisfying the following

DEFINITION 0.1 *Let E be a set and $I \subseteq E \times E$ an irreflexive and symmetric relation. A monoid given by a set of generators E and relations $ab = ba$ for all $(a, b) \in I$ is called free partially commutative and denoted by $M(E, I)$. If $(a, b) \in I$ then the members $a, b \in E$ are said to be commuting generators.*

This definition is varied with ordinary [4]. We do not demand that the set E is finite. It was shown in [12] that if the monoid $M(E, I)$ does not contain triples of mutually commuting generators, then its homological dimension ≤ 2 . We put forward a conjecture in [11] that if $M(E, I)$ does not contain n -tuples of mutually commuting generators, then its homological dimension $\leq n$. This conjecture was confirmed in the case of finite E by L. Polyakova [27]. We have confirmed it in the general case (Cor. 2.18).

We will prove that Leech homology groups of free partially commutative monoids are isomorphic to the cubical homology groups. This result we use to the building complexes for computing the homology groups of monoids and Hochschild homology of monoid rings.

Third section is devoted to the homology of $M(E, I)$ -sets. M. Bednarczyk [2] has introduced *asynchronous transition systems* to the modelling the concurrent processes. In [13] it was proved that the category of asynchronous transition systems admits an inclusion into the category of pointed sets over free partially commutative monoids. Thus asynchronous transition systems may be considered as $M(E, I)$ -sets. It allows us to build the functors from the category of asynchronous transition systems into the category of the abelian group and homomorphisms. Their values are homology groups of asynchronous transition systems. It was shown in [13] (see [12] as well) that these groups are isomorphic to the homology groups of $M(E, I)$ with coefficients in some right $M(E, I)$ -modules. Goubault [8] and Gaucher [6], [7] have given a definition of homology groups for higher dimensional automata which are precubical sets with additional structure. There is a functor from the category of asynchronous transition systems into the category of precubical sets. In the third section we prove that homology groups of asynchronous transition systems, defined in [13], are isomorphic to the homology groups of the corresponding precubical sets. In the work [12] it was proved that if an asynchronous transition system does not contain triples of independent events, then its homological dimension ≤ 2 . It was given a conjecture that if the asynchronous transition system does not contain n -tuples of independent

events, then its homological dimension $\leq n$ [12, Open Problem 2]. We have confirmed this conjecture.

We use the following notations:

Set	is the category of sets and maps.
Set _*	is the category of pointed sets: each its object is a set X with a selected element, written $*$ and called the “base point” ; its morphisms are maps $X \rightarrow Y$ which send the base point of X to that of Y . Such maps are called <i>based</i> . Following [30] we identify the category of sets and partial functions with Set_* .
Ab	is the category of abelian groups and homomorphisms.
$L : Set \rightarrow Ab$	is the left adjoint to the forgetful functor $U : Ab \rightarrow Set$; L assign to every set E the free abelian group $L(E)$ generated by E and to a map $f : E_1 \rightarrow E_2$ the canonical homomorphism $L(f) : L(E_1) \rightarrow L(E_2)$ such that $L(f)(e) = f(e)$ for all $e \in E_1$.
$\mathbb{I}$	is the totally ordered set $\{0, 1\}$ with the order relation $0 \leq 1$.
$\mathbb{Z}$	is the additive group of integers.
$\mathbb{N}$	is the set of non-negative integers or the additive monoid $\{0, 1, 2, 3, \dots\}$ or the free multiplicative monoid $\{1, a, a^2, a^3, \dots\}$.
$\mathbb{R}$	is the set of reals.

For any category $\mathcal{A}$, denote by $\mathcal{A}^{op}$ the dual category. For any pair $a, b \in Ob\mathcal{A}$, denote by $\mathcal{A}(a, b)$ the set of all morphisms $a \rightarrow b$. Given a small category $\mathcal{C}$ we denote by $\mathcal{A}^{\mathcal{C}}$ the category of functors $\mathcal{C} \rightarrow \mathcal{A}$ and natural transformations.

We will be consider any monoid as the small category with one object. This exert influence on our terminology. In particular a right M -set X will be considered and denoted as a functor $X : M^{op} \rightarrow Set$ (the value of X at the unique object will be denoted by $X(M)$ or shortly X). Morphisms of right M -sets are natural transformations.

1 Homology of categories and precubical sets

1.1 Precubical sets

A *precubical set* $X = (X_n, \partial_i^{n,\varepsilon})$ is a sequence of sets $(X_n)_{n \in \mathbb{N}}$ with a family of maps $\partial_i^{n,\varepsilon} : X_n \rightarrow X_{n-1}$, defined for $1 \leq i \leq n$, $\varepsilon \in \{0, 1\}$, for which the following diagrams is commutative for all $\alpha, \beta \in \{0, 1\}$, $n \geq 2$, $1 \leq i < j \leq n$:

$$\begin{array}{ccc} X_n & \xrightarrow{\partial_j^{n,\beta}} & X_{n-1} \\ \partial_i^{n,\alpha} \downarrow & & \downarrow \partial_i^{n-1,\alpha} \\ X_{n-1} & \xrightarrow{\partial_{j-1}^{n-1,\beta}} & X_{n-2} \end{array}$$

For example, if $X_n = \emptyset$ and $n > 1$, then the maps $\partial_1^{1,0} : X_1 \rightarrow X_0$ and $\partial_1^{1,1} : X_1 \rightarrow X_0$ will be determine a directed graph.

Precubical sets as functors. Let $\square_+$ be a category consisting of the finite sets $\mathbb{I}^n = \{0, 1\}^n$ ordered as the Cartesian power of $\mathbb{I}$. Any morphism of $\square_+$ is defined as an ascending map which admits a decomposition of the form $\delta_i^{k,\varepsilon} : \mathbb{I}^{k-1} \rightarrow \mathbb{I}^k$ where

$$\delta_i^{k,\varepsilon}(x_1, \dots, x_{k-1}) = (x_1, \dots, x_{i-1}, \varepsilon, x_i, \dots, x_{k-1}), \quad \varepsilon \in \mathbb{I}, 1 \leq i \leq k.$$

Every morphism $f \in \square_+(\mathbb{I}^m, \mathbb{I}^n)$ has a decomposition $f = \delta_{j_{n-m}}^{n,\varepsilon_{n-m}} \dots \delta_{j_1}^{m+1,\varepsilon_1}$, $1 \leq j_1 < \dots < j_{n-m} \leq n$ because of the equations $\delta_j^{n,\beta} \delta_i^{n-1,\alpha} = \delta_i^{n,\alpha} \delta_{j-1}^{n-1,\beta}$ for $1 \leq i < j \leq n$, $\alpha \in \{0, 1\}$, $\beta \in \{0, 1\}$. This allows us to define a precubical set as a functor $X : \square_+^{op} \rightarrow \text{Set}$ with values $X(\mathbb{I}^n) = X_n$ at $\mathbb{I}^n \in \text{Ob}(\square_+)$ and $X(f) = \partial_{j_1}^{m+1,\varepsilon_1} \dots \partial_{j_{n-m}}^{n,\varepsilon_{n-m}}$ at $f = \delta_{j_{n-m}}^{n,\varepsilon_{n-m}} \dots \delta_{j_1}^{m+1,\varepsilon_1} : \mathbb{I}^m \rightarrow \mathbb{I}^n$. A morphism of precubical sets is defined as a natural transformation. So we get the category of precubical sets $\text{Set}^{\square_+^{op}}$.

Integral homology groups of precubical sets.

DEFINITION 1.1 Let $X = (X_n, \partial_i^{n,\varepsilon})$ be a precubical set and $C_*(X)$ the chain complex of abelian groups $C_n(X) = L(X_n)$, $n \geq 0$, with $C_n(X) = 0$ for $n < 0$. Its differentials $d_n : C_n(X) \rightarrow C_{n-1}(X)$ are defined by

$$d_n = \sum_{i=1}^n (-1)^i (L(\partial_i^{n,1}) - L(\partial_i^{n,0})).$$

A n -th integral homology group of the precubical set X is the group $H_n(C_*(X))$.

Cubical subsets of Euclidian space. Consider precubical sets which integral homology groups are studied in [14]-[15].

For each integer $l \in \mathbb{Z}$ the closed intervals $[l, l+1] \subset \mathbb{R}$ and $\{l\} = [l, l] \subset \mathbb{R}$ are called *elementary intervals*. The elementary intervals $[l, l+1]$ are *nondegenerate* and $[l, l]$ *degenerate*.

DEFINITION 1.2 *An elementary cube is a Cartesian product of elementary intervals $Q = I_1 \times I_2 \times \cdots \times I_n \subset \mathbb{R}^n$. The number of nondegenerate intervals among $I_1, I_2, \dots, I_n$ is denoted by $\dim Q$. Denote the dimension of Euclidian space by $\text{emb } Q = n$.*

DEFINITION 1.3 *A cubical subset of Euclidian space $\mathbb{R}^n$ is the union of arbitrary elementary cubes $Q_1, Q_2, \dots$, for which $\text{emb } Q_i = n$.*

For any cubical subset $X \subseteq \mathbb{R}^n$ and $m \geq 0$ denote by $\mathcal{K}_m(X)$ the set of elementary cubes $Q \subseteq X$ such that $\dim Q = m$.

EXAMPLE 1.4 $X = [0, 1] \times [0, 2] \cup [1, 2] \times [0, 0] \cup [2, 2] \times [0, 1] \cup [1, 2] \times [1, 1]$ (*Pic. 1*).

$\mathcal{K}_0(X) = \{[0, 0] \times [0, 0], [1, 1] \times [0, 0], [2, 2] \times [0, 0], [0, 0] \times [1, 1], [1, 1] \times [1, 1], [2, 2] \times [1, 1], [0, 0] \times [2, 2], [1, 1] \times [2, 2]\}$.

$\mathcal{K}_1(X) = \{[0, 1] \times [0, 0], [1, 2] \times [0, 0], [0, 1] \times [1, 1], [1, 2] \times [1, 1], [0, 1] \times [2, 2], [0, 0] \times [0, 1], [1, 1] \times [0, 1], [2, 2] \times [0, 1], [0, 0] \times [1, 2], [1, 1] \times [1, 2]\}$.

$\mathcal{K}_2(X) = \{[0, 1] \times [0, 1], [0, 1] \times [1, 2]\}$.

Let $X \subseteq \mathbb{R}^n$ be a cubical subset. For any elementary cube $Q = I_1 \times I_2 \times \cdots \times I_n \in \mathcal{K}_m(X)$, where $I_{k_j} = [a_j, b_j]$ is a j -th nondegenerate elementary interval, we let

$$\partial_j^{m,0} Q = I_1 \times I_2 \times \cdots \times I_{k_j-1} \times \{a_j\} \times I_{k_j+1} \times \cdots \times I_n \quad ,$$

$$\partial_j^{m,1} Q = I_1 \times I_2 \times \cdots \times I_{k_j-1} \times \{b_j\} \times I_{k_j+1} \times \cdots \times I_n \quad ,$$

where $1 \leq j \leq \dim Q = m$. We have got the precubical set $\mathcal{K}(X) = (\mathcal{K}_m(X), \partial_j^{m,\varepsilon})$. Consequently, the cubical subsets of Euclidian spaces may be regarded as precubical sets. Easy to see (from [15, Corollary 2.3], for example), that the homology groups of precubical sets corresponding to the cubical subsets of Euclidian spaces are isomorphic to the cubical homology groups defined in [15]. In particular it is true the following assertion:




Figure 1: A cubical subset of the space $\mathbb{R}^2$.

Lemma 1.1 *Let $C_*(\mathcal{K}([0, 1]^n))$ be a chain complex of abelian groups $C_m(\mathcal{K}([0, 1]^n)) = L(\mathcal{K}_m([0, 1]^n))$ generated by cubes $Q \subseteq [0, 1]^n$ for which $\dim Q = m$. Let $Q = I_1 \times \cdots \times I_n$ be an elementary cube which equals a product of m elementary nondegenerate intervals $I_{k_1} = \cdots = I_{k_m} = [0, 1]$, $1 \leq k_1 < k_2 < \cdots < k_m \leq n$, and $n - m$ degenerate elementary intervals. Define the differentials d_m of complex $C_*(\mathcal{K}([0, 1]^n))$ as*

$$d_m Q = \sum_{j=1}^m (-1)^j (L(\partial_j^{m,1})Q - L(\partial_j^{m,0})Q).$$

Then

$$H_q(C_*(\mathcal{K}([0, 1]^n))) = \begin{cases} \mathbb{Z}, & \text{if } q = 0, \\ 0, & \text{if } q > 0. \end{cases}$$

This assertion is proved in [14, Prop. 4.61] for the most general case where an arbitrary elementary cube of Euclidian space is taken instead of $[0, 1]^n$.

1.2 Smith normal form and calculating the homology groups

Let $G' \xrightarrow{\alpha} G \xrightarrow{\beta} G''$ be homomorphisms of abelian groups such that $\beta \circ \alpha = 0$. Then we can consider the homology group $\text{Ker}(\beta)/\text{Im}(\alpha)$. Let us describe

a method of calculation for this homology group when G' , G , and G'' are finitely generated free abelian groups.

Proposition 1.2 [24, Theorem III.4(43)] *Let*

$$A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{pmatrix}$$

be a matrix with integer entries $a_{ij} \in \mathbb{Z}$. Then there is an $m \times m$ matrix T and an $n \times n$ matrix S with integer entries such that:

- (i) $\det(T) = \pm 1$, $\det(S) = \pm 1$;
- (ii) $A = T \circ D(A) \circ S$ for a natural number $k \geq 0$ and $m \times n$ matrix

$$D(A) = \begin{pmatrix} d_1 & 0 & \cdots & 0 & 0 & \cdots & 0 \\ 0 & d_2 & \cdots & 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & d_k & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 & 0 & \cdots & 0 \\ \vdots & \vdots & & \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & 0 & 0 & \cdots & 0 \end{pmatrix}$$

all entries of which are equal to 0 except the diagonal numbers $d_1 \leq d_2 \leq \cdots \leq d_k$ which satisfy that d_i divides d_{i+1} for all $1 \leq i \leq k-1$.

The matrix $D(A)$ is said to be a *Smith normal form* of A . This form is used for the computation of the homology groups of simplicial complexes in [29]. We refer the reader to [9], where an algorithm is presented for computing the Smith normal form of an integer matrix, which performs well in practice. There are packages such as GAP for the computation of the Smith normal form.

Proposition 1.3 *Let a homomorphism $\alpha : \mathbb{Z}^n \rightarrow \mathbb{Z}^m$ be given by a $m \times n$ matrix A and $\beta : \mathbb{Z}^m \rightarrow \mathbb{Z}^p$ by $p \times m$ matrix B . Suppose the sequence $\mathbb{Z}^n \xrightarrow{\alpha} \mathbb{Z}^m \xrightarrow{\beta} \mathbb{Z}^p$ satisfies to $\beta \circ \alpha = 0$. If $d_1, d_2, \dots, d_k$ are the nonzero diagonal entries of the Smith normal form $D(A)$, then*

$$\text{Ker } \beta / \text{Im } \alpha \cong \mathbb{Z}/d_1\mathbb{Z} \times \mathbb{Z}/d_2\mathbb{Z} \times \cdots \times \mathbb{Z}/d_k\mathbb{Z} \times \mathbb{Z}^{m-k-b} \quad (1)$$

where b is the rank of the matrix B and may be computed as the number of nonzero entries of $D(B)$.

1.3 Homology of small categories

In this section it is considered homology groups of small categories $\mathcal{C}$ with coefficients in functors $F : \mathcal{C} \rightarrow \text{Ab}$.

Homology of categories and derived functors of the colimit.

DEFINITION 1.5 Homology groups of a small category $\mathcal{C}$ with coefficients in a functor $F : \mathcal{C} \rightarrow \text{Ab}$ are homology groups of the chain complex $C_*(\mathcal{C}, F)$ consisting of the abelian groups

$$C_n(\mathcal{C}, F) = \bigoplus_{c_0 \rightarrow \cdots \rightarrow c_n} F(c_0), \quad n \geq 0,$$

with differentials $d_n = \sum_{i=0}^n (-1)^i d_i^n : C_n(\mathcal{C}, F) \rightarrow C_{n-1}(\mathcal{C}, F)$, $n > 0$, where for each

$$(c_0 \xrightarrow{\alpha_1} c_1 \xrightarrow{\alpha_2} \cdots \xrightarrow{\alpha_n} c_n, a) \in \bigoplus_{c_0 \rightarrow \cdots \rightarrow c_n} F(c_0), \quad a \in F(c_0)$$

d_i^n are defined by

$$d_i^n(c_0 \rightarrow \cdots \rightarrow c_n, a) = \begin{cases} (c_0 \rightarrow \cdots \rightarrow \widehat{c_i} \rightarrow \cdots \rightarrow c_n, a) & , \text{ for } 1 \leq i \leq n \\ (c_1 \rightarrow \cdots \rightarrow c_n, F(c_0 \xrightarrow{\alpha_1} c_1)(a)) & , \text{ for } i = 0. \end{cases}$$

It is well known [5, Prop. 3.3, Application 2] that there exists an isomorphism of left satellites of the colimit $\varinjlim^{\mathcal{C}} : \text{Ab}^{\mathcal{C}} \rightarrow \text{Ab}$ and the functors $H_n(C_*(\mathcal{C}, -)) : \text{Ab}^{\mathcal{C}} \rightarrow \text{Ab}$. Since the category $\text{Ab}^{\mathcal{C}}$ has enough projectives, these satellites are natural isomorphic to the left derived functor of $\varinjlim^{\mathcal{C}} : \text{Ab}^{\mathcal{C}} \rightarrow \text{Ab}$. Denote the values $H_n(C_*(\mathcal{C}, F))$ of the satellites at $F \in \text{Ab}^{\mathcal{C}}$ by $\varinjlim_n^{\mathcal{C}} F$.

Let $\mathcal{C}$ be a small category. Denote by $\Delta_{\mathcal{C}} \mathbb{Z} : \mathcal{C} \rightarrow \text{Ab}$, or shortly $\Delta \mathbb{Z}$, the functor which has constant values $\Delta \mathbb{Z}(c) = \mathbb{Z}$ at $c \in \text{Ob}(\mathcal{C})$ and $\Delta \mathbb{Z}(\alpha) = 1_{\mathbb{Z}}$ at $\alpha \in \text{Mor}(\mathcal{C})$. Integral homology groups of $\mathcal{C}$ are the groups $\varinjlim_n^{\mathcal{C}} \Delta_{\mathcal{C}} \mathbb{Z}$.

EXAMPLE 1.6 Let $pt = [0]$ be a category consisting of one object 0 and one morphism 1_0 . Then for every $n \geq 0$ the abelian group $C_n(pt, \Delta_{pt} \mathbb{Z})$ is generated by the element e_n which is equal to the sequence of n identity morphisms $0 \xrightarrow{1_0} 0 \xrightarrow{1_0} \cdots \xrightarrow{1_0} 0$. Since $d_i^n(e_n) = e_{n-1}$,

$$d_n(e_n) = \begin{cases} e_{n-1}, & \text{if } n \text{ is even,} \\ 0, & \text{if } n \text{ is odd.} \end{cases}$$

Consequently the chain $C_*(\mathcal{C}, \Delta \mathbb{Z})$ is isomorphic to the chain

$$0 \leftarrow \mathbb{Z} \xleftarrow{0} \mathbb{Z} \xleftarrow{1} \mathbb{Z} \xleftarrow{0} \mathbb{Z} \xleftarrow{1} \dots,$$

which n -th homology groups are equal to 0 if $n > 0$, and $H_0(C_*(pt, \Delta \mathbb{Z})) \cong \mathbb{Z}$. It follows that

$$\varinjlim_n^{pt} \Delta \mathbb{Z} = \begin{cases} \mathbb{Z}, & \text{if } n = 0, \\ 0, & \text{if } n > 0. \end{cases}$$

Lemma 1.4 *For arbitrary small category $\mathcal{C}$ and all $n \geq 0$ there are isomorphisms $\varinjlim_n^{C^{op}} \Delta_{C^{op}} \mathbb{Z} \cong \varinjlim_n^{\mathcal{C}} \Delta_{\mathcal{C}} \mathbb{Z}$.*

PROOF. Consider the homomorphisms $C_n(\mathcal{C}^{op}, \Delta \mathbb{Z}) \rightarrow C_n(\mathcal{C}, \Delta \mathbb{Z})$, which assign to any element $(c_0 \xrightarrow{\alpha_1^{op}} c_1 \rightarrow \dots \xrightarrow{\alpha_n^{op}} c_n)$ of $C_n(\mathcal{C}^{op}, \Delta \mathbb{Z})$ the element $(c_0 \xleftarrow{\alpha_1} c_1 \leftarrow \dots \xleftarrow{\alpha_n} c_n)$ for even n , and $-(c_0 \xleftarrow{\alpha_1} c_1 \leftarrow \dots \xleftarrow{\alpha_n} c_n)$ for odd n . These homomorphisms are isomorphisms $C_*(\mathcal{C}^{op}, \Delta \mathbb{Z}) \xrightarrow{\cong} C_*(\mathcal{C}, \Delta \mathbb{Z})$. Consequently the homology groups of the complexes are isomorphic. $\square$

Tensor product of functors. Let $Lh^c : \text{Ab}^{\mathcal{C}} \rightarrow \text{Ab}$ be the composition of the functors $L : \text{Set} \rightarrow \text{Ab}$ and $h^c(-) = \mathcal{C}(c, -) : \mathcal{C} \rightarrow \text{Set}$. For a functor $F : \mathcal{C}^{op} \rightarrow \mathcal{A}$ and an object $A \in \mathcal{A}$ we denote by $\text{Hom}(F, A)$ the functor $\mathcal{C} \rightarrow \text{Ab}$, having the values $\mathcal{A}(F(c), A)$ at $c \in \text{Ob}(\mathcal{C})$ and assigning to every $\alpha \in \mathcal{C}(c_1, c_2)$ the map $\text{Hom}(F, A)(\alpha) : \mathcal{A}(F(c_2), A) \rightarrow \mathcal{A}(F(c_1), A)$ defined as $\text{Hom}(F, A)(\alpha)(f) = f \circ F(\alpha)$.

Lemma 1.5 *Let $\mathcal{C}$ be a small category and $\mathcal{A}$ an additive category with infinite coproducts. Then there exists a bifunctor $\otimes : \text{Ab}^{\mathcal{C}} \times \mathcal{A}^{C^{op}} \rightarrow \mathcal{A}$ which is additive at each argument and has the following properties*

- *there are isomorphisms*

$$\mathcal{A}(G \otimes F, A) \xrightarrow{\cong} \text{Ab}^{\mathcal{C}}(G, \text{Hom}(F, A))$$

natural in $G \in \text{Ab}^{\mathcal{C}}$, $F \in \mathcal{A}^{C^{op}}$, and $A \in \mathcal{A}$;

- *there are isomorphisms*

$$Lh^c \otimes F \xrightarrow{\cong} F(c)$$

natural at $c \in \mathcal{C}$ $F \in \mathcal{A}^{C^{op}}$;

- for each $F \in \mathcal{A}^{C^{op}}$ the functor $(-) \otimes F : \text{Ab}^{\mathcal{C}} \rightarrow \mathcal{A}$ preserves the colimits;
- for every $G \in \text{Ab}^{\mathcal{C}}$ the functor $G \otimes (-) : \mathcal{A}^{C^{op}} \rightarrow \mathcal{A}$ preserves the colimits.

PROOF. Let $\mathcal{B}$ and $\mathcal{A}$ be additive categories. Denote the category of additive functors $\mathcal{B} \rightarrow \mathcal{A}$ by $\text{Add}(\mathcal{B}, \mathcal{A})$. It is well known [23, p.11], that for each small category $\mathcal{C}$ there is the additive category $\text{Mat } \mathbb{Z}\mathcal{C}$ which contains $\mathcal{C}$ and has the isomorphisms $\mathcal{A}^{\mathcal{C}} \cong \text{Add}(\text{Mat } \mathbb{Z}\mathcal{C}, \mathcal{A})$ for every additive category $\mathcal{A}$. There is a bifunctor [28, Theorem 17.7.2]

$$\otimes_{C^{op}} : \text{Add}(\text{Mat } \mathbb{Z}\mathcal{C}, \text{Ab}) \times \text{Add}(\text{Mat } \mathbb{Z}C^{op}, \mathcal{A}) \rightarrow \mathcal{A},$$

for which firstly there are natural isomorphisms $h^c \otimes F \cong F(c)$, and secondly the functors $(-) \otimes F$ and $G \otimes (-)$ preserves the colimits. Using this bifunctor and the isomorphisms of categories

$$\text{Add}(\text{Mat } \mathbb{Z}\mathcal{C}, \text{Ab}) \cong \text{Ab}^{\mathcal{C}}, \quad \text{Add}(\text{Mat } \mathbb{Z}C^{op}, \mathcal{A}) \cong \mathcal{A}^{C^{op}}$$

we obtain the bifunctor $\otimes : \text{Ab}^{\mathcal{C}} \times \mathcal{A}^{C^{op}} \rightarrow \mathcal{A}$ with required properties. $\square$

Derived functors of the colimit and partial derived of the tensor product at the first variable. Let $\mathcal{C}$ be a small category. Recall that the homology groups of the category C^{op} is denoted by $\varinjlim_n^{C^{op}} : \text{Ab}^{C^{op}} \rightarrow \text{Ab}$. The following lemma is well known [25] and it may be proved from general approach. We prove it by the definition of $\varinjlim_n^{C^{op}}$ as the homology groups of the chain complex $C_*(\mathcal{C}, F)$.

Lemma 1.6 *For every projective resolution*

$$0 \leftarrow \Delta_{\mathcal{C}} \mathbb{Z} \xleftarrow{\varepsilon} P_0 \xleftarrow{d_1} P_1 \leftarrow \dots \xleftarrow{d_n} P_n \leftarrow \dots$$

the n -th homology group of the chain complex $P_* \otimes F$ is isomorphic to $\varinjlim_n^{C^{op}} F$.

PROOF. Let $P_*(\mathcal{C})$ be the chain complex in the category $\text{Ab}^{\mathcal{C}}$ where

$$P_n(\mathcal{C}) = \bigoplus_{c_0 \rightarrow \dots \rightarrow c_n} Lh^{c_n} \in \text{Ab}^{\mathcal{C}}$$

are projective objects in $\text{Ab}^{\mathcal{C}}$. The morphisms $d_n : P_n(\mathcal{C}) \rightarrow P_{n-1}(\mathcal{C})$ are defined for $n > 0$ as $\sum_{i=0}^n (-1)^i d_i^n$ where $d_i^n : P_n(\mathcal{C}) \rightarrow P_{n-1}(\mathcal{C})$ are natural

transformations which components $(d_i^n)_c$ for $c \in \mathcal{C}$ are given on generators of $\bigoplus_{c_0 \rightarrow \dots \rightarrow c_n} L\mathcal{C}(c_n, c)$ as $(d_i^n)_c(c_0 \rightarrow \dots \rightarrow c_n \rightarrow c) = (c_0 \rightarrow \dots \rightarrow \widehat{c_i} \rightarrow \dots \rightarrow c_n \rightarrow c)$. Define a natural transformation $d_0 : P_0(\mathcal{C}) \rightarrow \Delta_{\mathcal{C}}\mathbb{Z}$ by letting $(d_0)_c(c_0 \rightarrow c) = c$ on generators of the group $P_0(\mathcal{C})(c)$ where $\Delta_{\mathcal{C}}\mathbb{Z}(c)$ is considered as the free abelian group with one generator c . We will prove that the sequence

$$0 \rightarrow \Delta_{\mathcal{C}}\mathbb{Z} \xleftarrow{d_0} P_0(\mathcal{C}) \xleftarrow{d_1} P_1(\mathcal{C}) \leftarrow \dots \xleftarrow{d_n} P_n(\mathcal{C}) \leftarrow \dots$$

is exact in $\text{Ab}^{\mathcal{C}}$. Denote this sequence by $\overline{P}_*(\mathcal{C})$. Let $s_n : \overline{P}_{n-1}(\mathcal{C})(c) \rightarrow \overline{P}_n(\mathcal{C})(c)$ be homomorphisms defined for $n > 0$ on generators by

$$s_n(c_0 \rightarrow \dots \rightarrow c_{n-1} \rightarrow c) = (c_0 \rightarrow \dots \rightarrow c_{n-1} \rightarrow c \xrightarrow{1_c} c),$$

and for $n = 0$ by $s_0(c) = (c \xrightarrow{1_c} c)$. It is easy to see that s_n make up the contracting homotopy of the chain complex $\overline{P}_*(\mathcal{C})(c)$. Hence the sequence $\overline{P}_*(\mathcal{C})$ is exact in $\text{Ab}^{\mathcal{C}}$. Thus $P_*(\mathcal{C})$ is the projective resolution of $\Delta_{\mathcal{C}}\mathbb{Z}$. It is easy to see from Lemma 1.5 that the chain complex

$$0 \leftarrow \Delta_{\mathcal{C}}\mathbb{Z} \otimes F \xleftarrow{d_0 \otimes F} P_0(\mathcal{C}) \otimes F \leftarrow \dots \leftarrow P_{n-1}(\mathcal{C}) \otimes F \xleftarrow{d_n \otimes F} P_n(\mathcal{C}) \otimes F \leftarrow \dots$$

is isomorphic to $C_*(\mathcal{C}^{op}, F)$. Consequently $H_n(P_*(\mathcal{C}) \otimes F) \cong \varinjlim_n^{\mathcal{C}^{op}} F$.

Now consider an arbitrary projective resolution P_* of the object $\Delta_{\mathcal{C}}\mathbb{Z}$ in $\text{Ab}^{\mathcal{C}}$. It is homotopical equivalent to $P_*(\mathcal{C})$. Since the functor $(-) \otimes F$ is additive, it sends homotopical equivalent chain complexes to homotopical equivalent chain complexes. Consequently $H_n(P_* \otimes F) \cong H_n(P_*(\mathcal{C}) \otimes F) \cong \varinjlim_n^{\mathcal{C}^{op}} F$. $\square$

Coinitial functors. A small category $\mathcal{C}$ is *connected* if for any objects $a, b \in \mathcal{C}$ there exists a sequence of morphisms $a = c_0 \rightarrow c_1 \leftarrow c_2 \rightarrow \dots \leftarrow c_{2k} = b$. $\mathcal{C}$ is *acyclic* if $\varinjlim_n^{\mathcal{C}} \Delta\mathbb{Z} = 0$ for all $n > 0$. Let $S : \mathcal{C} \rightarrow \mathcal{D}$ be a functor from a small category into an arbitrary category. For any $d \in \text{Ob}(\mathcal{D})$ a *fibre* (or a *comma-category*) S/d is the category which objects are pairs (c, α) , consisting of objects $c \in \mathcal{C}$ and morphisms $\alpha \in \mathcal{D}(S(c), d)$. Morphisms in S/d are defined as triples (f, α_1, α_2) of $f \in \mathcal{C}(c_1, c_2)$, $\alpha_1 \in \mathcal{D}(S(c_1), d)$, $\alpha_2 \in \mathcal{D}(S(c_2), d)$ for which $\alpha_2 \circ S(f) = \alpha_1$. A *forgetful functor* $Q_d : S/d \rightarrow \mathcal{C}$ of the fibre is defined as $Q_d(c, \alpha) = c$ on objects, and $Q_d(f, \alpha_1, \alpha_2) = f$ at morphisms. If S is a full embedding $\mathcal{C} \subseteq \mathcal{D}$, then S/d is denoted by $\mathcal{C}/d$.

DEFINITION 1.7 *A functor $S : \mathcal{C} \rightarrow \mathcal{D}$ between small categories is strong coinital if for all object $d \in \mathcal{D}$ the categories S/d are connected and acyclic.*

This is holds if and only if for all $d \in \mathcal{D}$

$$\varinjlim_n^{S/d} \Delta \mathbb{Z} = \begin{cases} 0, & \text{if } n > 0 \\ \mathbb{Z}, & \text{if } n = 0. \end{cases}$$

A cofibre d/S of an object $d \in \mathcal{D}$ over $S : \mathcal{C} \rightarrow \mathcal{D}$ is a category with objects (c, α) where $c \in \text{Ob}(\mathcal{C})$, $\alpha \in \mathcal{D}(d, S(c))$. Its morphisms $(c_1, \alpha_1) \rightarrow (c_2, \alpha_2)$ are given by triples (f, α_1, α_2) of morphisms $f \in \mathcal{C}(c_1, c_2)$, $\alpha_1 \in \mathcal{D}(d, S(c_1))$, $\alpha_2 \in \mathcal{D}(d, S(c_2))$ satisfying $S(f) \circ \alpha_1 = \alpha_2$.

Proposition 1.7 *Let $\mathcal{C}$ and $\mathcal{D}$ be small categories and $S : \mathcal{C} \rightarrow \mathcal{D}$ a functor. Then there are canonical homomorphisms $\varinjlim_n^{C^{op}} (F \circ S^{op}) \rightarrow \varinjlim_n^{D^{op}} F$ which are natural in F . These homomorphisms are isomorphisms for all $n \geq 0$ if and only if the functor S is strong coinital.*

PROOF. This follows from the assertion [26, Theorem 2.3] affirmed for homomorphisms $\varinjlim_n^{\mathcal{C}} (F \circ S) \rightarrow \varinjlim_n^{\mathcal{D}} F$. In this case we recive the following necessary and sufficient condition $\varinjlim_n^{d/S} \Delta \mathbb{Z} \cong \varinjlim_n^{pt} \Delta \mathbb{Z}$ under which these homomorphisms are isomorphisms. We substitute S by S^{op} and obtain that S is strong coinital if and only if $\varinjlim_n^{(S/d)^{op}} \Delta \mathbb{Z} \cong \varinjlim_n^{pt} \Delta \mathbb{Z}$. But $\varinjlim_n^{(S/d)^{op}} \Delta \mathbb{Z} \cong \varinjlim_n^{S/d} \Delta \mathbb{Z}$ by Lemma 1.4. $\square$

Corollary 1.8 *The composition of strong coinital functors is strong coinital.*

Homology of $\mathcal{D}$ —sets. Let $S : \mathcal{C} \rightarrow \mathcal{D}$ be a functor between small categories. Then for a cocomplete category $\mathcal{A}$ the functor $(-) \circ S : \mathcal{A}^{\mathcal{D}} \rightarrow \mathcal{A}^{\mathcal{C}}$ has (see [20]) a left adjoint functor Lan^S which value at $F \in \mathcal{A}^{\mathcal{C}}$ is called by a *left Kan extension* $\text{Lan}^S F$ of $F \in \mathcal{A}^{\mathcal{C}}$ along S . A *connection components of a category* are its maximal connected subcategories. Let $\mathcal{C}$ be a small category in which every connection component has an initial object. For each object $c \in \mathcal{C}$ we select one initial object $i(c)$ in the connection component contained c . Denote by $i_c : i(c) \rightarrow c$ the unique morphism from $i(c)$ into c . Let $\text{init}(\mathcal{C})$ be the set consisting of the selected initial objects.

Lemma 1.9 *Let $\mathcal{A}$ be a cocomplete category. Suppose that $S : \mathcal{C} \rightarrow \mathcal{D}$ is a functor between small categories such that for each object $d \in \mathcal{D}$ every connection component of d/S has an initial object. Then for any functor $F : \mathcal{C}^{op} \rightarrow \mathcal{A}$ the left Kan extension $\text{Lan}^{S^{op}} F : \mathcal{D}^{op} \rightarrow \mathcal{A}$ is isomorphic to a functor which values on objects $d \in \mathcal{D}$ equal $\coprod_{\beta \in \text{init}(d/S)} FQ_d^{op}(\beta)$ and assigned to every morphism $\alpha : d \rightarrow e$ of the category $\mathcal{D}$ the morphism $\overline{\alpha}$ of $\mathcal{A}$ determined by commutativity $\forall \gamma \in \text{init}(e/S)$ of the following diagrams*

$$\begin{array}{ccc}
\coprod_{\gamma \in \text{init}(e/S)} FQ_e^{op}(\gamma) & \xrightarrow{\overline{\alpha}} & \coprod_{\beta \in \text{init}(d/S)} FQ_d^{op}(\beta) \\
\uparrow \text{in}_\gamma & & \uparrow \text{in}_{i(\gamma \circ \alpha)} \\
FQ_e^{op}(\gamma) = FQ_d^{op}(\gamma \circ \alpha) & \xrightarrow{FQ_d^{op}(i_{\gamma \circ \alpha})} & FQ_d^{op}(i(\gamma \circ \alpha))
\end{array}$$

PROOF. Substitute the category $\mathcal{A}$ by $\mathcal{A}^{op}$ in [18, 1.1]. Since products in the dual category are coproducts and arrows have the dual direction, the assertion follows from [18, 1.1]. $\square$

Let $\mathcal{D}$ be a small category and $X \in \text{Set}^{\mathcal{D}^{op}}$ a functor. Denote by $\mathcal{D}/X$ a small category which objects are members $x \in \coprod_{d \in \text{Ob } \mathcal{D}} X(d)$ and its morphisms are triples (α, x, x') , with $\alpha \in \mathcal{D}(d, d')$, $x \in X(d)$, and $x' \in X(d')$ satisfying to $X(\alpha)(x') = x$.

It is also concerned with $\mathcal{D}$ and X the fibre h_*/X of Yoneda embedding $h_* : \mathcal{D} \rightarrow \text{Set}^{\mathcal{D}^{op}}$. Since h_* is a full embedding, every object $(d, \eta : h_d \rightarrow X)$ is determinant by the second element $\eta : h_d \rightarrow X$. Thus h_*/X may be considered as the category with objects $\eta : h_d \rightarrow X$ and morphisms given by commutative triangles

$$\begin{array}{ccc}
h_d & \xrightarrow{\eta} & X \\
& \searrow h_\alpha & \nearrow \eta' \\
& h_{d'} &
\end{array}$$

It is well known [20] that there is an isomorphism $\mathcal{D}/X \xrightarrow{\cong} h_*/X$. It is built on objects as the inverse to the map $(\eta : h_d \rightarrow X) \mapsto \eta_d(1_d)$. The value of the map at x is denoted by $\tilde{x} : h_d \rightarrow X$.

Let $\mathcal{D}$ be a small category and $X \in \text{Set}^{\mathcal{D}^{op}}$ and $F : (\mathcal{D}/X)^{op} \rightarrow \text{Ab}$ be functors.

Proposition 1.10 *For all $n \geq 0$ there are isomorphisms $\varinjlim_n^{(\mathcal{D}/X)^{op}} F \cong \varinjlim_n^{\mathcal{D}^{op}} \check{F}$ where $\check{F}(d) = \bigoplus_{x \in X(d)} F(x)$ and $\check{F}(\alpha) : \check{F}(a) \rightarrow \check{F}(b)$ at $\alpha \in \mathcal{D}(a, b)$ are morphisms making the following diagrams commutative*

$$\begin{array}{ccc}
\bigoplus_{x \in X(a)} F(x) & \xrightarrow{\check{F}(\alpha)} & \bigoplus_{x \in X(b)} F(x) \\
\uparrow \text{in}_x & & \text{in}_{X(\alpha)(x)} \uparrow \\
F(x) & \xrightarrow{F(X(\alpha)(x) \xrightarrow{\alpha} x)} & F(X(\alpha)(x))
\end{array} \tag{2}$$

PROOF. Denote $\check{F} = \text{Lan}^{Q^{op}} F$ where $Q : h_*/X \rightarrow \mathcal{D}$ is the forgetful full defined by the assignments $\tilde{x} : h_d \rightarrow X \mapsto d \in \mathcal{D}$ and $(\alpha, x, y) \mapsto \alpha$. For any $a \in \mathcal{D}$ each connected component of the category a/Q has an initial object. The values of the map $i : \text{Ob}(a/Q) \rightarrow \text{init}(a/Q)$ at $(\tilde{x} : h_c \rightarrow X, \alpha : a \rightarrow c)$ equal $i(\tilde{x}, \alpha) = (\tilde{x} \circ h_\alpha, 1_a)$ and $i_{(\tilde{x}, \alpha)} = (\alpha, \tilde{x} \circ h_\alpha, \tilde{x})$. Since $\tilde{x} \circ h_\alpha = \widetilde{X(\alpha)(x)}$ and the categories $\mathcal{D}/X$ and h_*/X may be identified, we obtain from Lemma 1.9 a definition of homomorphisms $\check{F}(\alpha)$ as making squares (2) commutative.

Since the coproduct functor is exact in Ab , $\text{Lan}^{Q^{op}}$ is exact. Consider an arbitrary projective resolution of an object $F \in \text{Ab}^{(\mathcal{D}/X)^{op}}$

$$0 \leftarrow F \leftarrow P_0 \leftarrow P_1 \leftarrow \dots$$

The functor $\text{Lan}^{Q^{op}}$ is left adjoint to the exact functor $(-) \circ Q^{op} : \text{Ab}^{\mathcal{D}^{op}} \rightarrow \text{Ab}^{(\mathcal{D}/X)^{op}}$. In accordance with a dual assertion to [3, Prop. 6.3] it carries projective objects to projective. Consequently it carries the projective resolution of F to the projective resolution of $\text{Lan}^{Q^{op}} F$. Applying the functor $\varinjlim^{\mathcal{D}^{op}}$ to the obtained resolution we get a sequence

$$0 \leftarrow \varinjlim^{(\mathcal{D}/X)^{op}} \text{Lan}^{Q^{op}} F \leftarrow \varinjlim^{(\mathcal{D}/X)^{op}} \text{Lan}^{Q^{op}} P_0 \leftarrow \varinjlim^{(\mathcal{D}/X)^{op}} \text{Lan}^{Q^{op}} P_1 \leftarrow \dots$$

such that homology groups of the chain complex

$$0 \leftarrow \varinjlim^{(\mathcal{D}/X)^{op}} \text{Lan}^{Q^{op}} P_0 \leftarrow \varinjlim^{(\mathcal{D}/X)^{op}} \text{Lan}^{Q^{op}} P_1 \leftarrow \dots$$

are isomorphic to groups $\varinjlim_n^{(\mathcal{D}/X)^{op}} \text{Lan}^{Q^{op}} F$. On the other hand there is the isomorphism $\varinjlim^{(\mathcal{D}/X)^{op}} \text{Lan}^{Q^{op}} \cong \varinjlim^{\mathcal{D}^{op}}$. Thus the homology groups of the chain complex are isomorphic to $\varinjlim_n^{\mathcal{D}^{op}} F$. $\square$

1.4 Homology of precubical sets

DEFINITION 1.8 A homological system of abelian groups on a precubical set $X \in \text{Set}^{\square_+^{op}}$ is an arbitrary functor $F : (\square_+/X)^{op} \rightarrow \text{Ab}$.

We will consider values of the functor

$$\varinjlim_n^{(\square_+/X)^{op}} : \text{Ab}^{(\square_+/X)^{op}} \rightarrow \text{Ab},$$

at homological systems of abelian groups on precubical sets $X : \square_+^{op} \rightarrow \text{Set}$.

Homology of a precubical abelian group.

Lemma 1.11 Let C_* be a chain complex

$$0 \leftarrow L\square_+(\mathbb{I}^0, \mathbb{I}^n) \xleftarrow{d_1} L\square_+(\mathbb{I}^1, \mathbb{I}^n) \xleftarrow{d_2} \dots \xleftarrow{d_n} L\square_+(\mathbb{I}^n, \mathbb{I}^n) \leftarrow 0,$$

consisting of free abelian groups C_k generated by morphisms $\mathbb{I}^k \rightarrow \mathbb{I}^n$. Its differentials are defined as $d_k = \sum_{i=1}^k (-1)^i (L(\partial_i^{k,1}) - L(\partial_i^{k,0}))$ where $\partial_i^{k,\varepsilon}(\sigma) = \square_+(\delta_i^{k,\varepsilon}, \mathbb{I}^n)(\sigma) = \sigma \circ \delta_i^{k,\varepsilon}$. Then its homology groups $H_q(C_*)$ are equal to 0 if $q > 0$ and $H_0(C_*) = \mathbb{Z}$.

PROOF. For $1 \leq i \leq j \leq k-1$ there is equations $\delta_{j+1}^{k,\beta} \delta_i^{k-1,\alpha} = \delta_i^{k,\alpha} \delta_j^{k-1,\beta}$. Therefore each morphism $\sigma : \mathbb{I}^k \rightarrow \mathbb{I}^n$ has an unique decomposition

$$\mathbb{I}^k \xrightarrow{\delta_1^{k+1,\varepsilon_1}} \mathbb{I}^{k+1} \xrightarrow{\delta_2^{k+2,\varepsilon_2}} \mathbb{I}^{k+2} \rightarrow \dots \xrightarrow{\delta_{j_1-1}^{k+j_1-1,\varepsilon_{j_1-1}}} \mathbb{I}^{k+j_1-1} \xrightarrow{\delta_{j_1+1}^{k+j_1,\varepsilon_{j_1+1}}} \dots \xrightarrow{\delta_{n-k}^{n,\varepsilon_{n-k}}} \mathbb{I}^n,$$

which may be written as

$$\sigma = \delta_{n-k}^{n,\varepsilon_{n-k}} \dots \delta_{j_1+1}^{k+j_1,\varepsilon_{j_1+1}} \delta_{j_1-1}^{k+j_1-1,\varepsilon_{j_1-1}} \dots \delta_2^{k+2,\varepsilon_2} \delta_1^{k+1,\varepsilon_1}.$$

To the studying the action of $\partial_i^{k,\varepsilon}$ at σ , $\partial_i^{k,\varepsilon}(\sigma) = \sigma \circ \delta_i^{k,\varepsilon}$, we denote morphisms $\delta_i^{k,\varepsilon}$ shortly by δ_i^ε . Consider a diagram:

$$\begin{array}{ccccccc} \mathbb{I}^k & \xrightarrow{\delta_1^\varepsilon} & \mathbb{I}^{k+1} & \xrightarrow{\delta_2^\varepsilon} \dots \xrightarrow{\delta_{j_1-1}^{\varepsilon_{j_1-1}}} & \mathbb{I}^{k+j_1-1} & \xrightarrow{\delta_{j_1+1}^{\varepsilon_{j_1+1}}} \dots \xrightarrow{\delta_{j_i-1}^{\varepsilon_{j_i-1}}} & \mathbb{I}^{k+j_i} \xrightarrow{\delta_{j_i+1}^{\varepsilon_{j_i+1}}} \dots \xrightarrow{\delta_{n-k}^{\varepsilon_{n-k}}} \mathbb{I}^n \\ \uparrow \delta_i^\varepsilon & & \uparrow \delta_{i+1}^\varepsilon & & \uparrow \delta_{i+j_1-1}^\varepsilon & & \uparrow \delta_{i+j_i-i}^\varepsilon \\ \mathbb{I}^{k-1} & \xrightarrow{\delta_1^\varepsilon} & \mathbb{I}^k & \xrightarrow{\delta_2^\varepsilon} \dots \xrightarrow{\delta_{j_1-1}^{\varepsilon_{j_1-1}}} & \mathbb{I}^{k+j_1-2} & \xrightarrow{\delta_{j_1+1}^{\varepsilon_{j_1+1}}} \dots \xrightarrow{\delta_{j_i-1}^{\varepsilon_{j_i-1}}} & \mathbb{I}^{k+j_i-1} \end{array}$$

The lower indexes of the morphism δ increase by 1 at the horizontal arrows except the arrows in the places $j_1, j_2, \dots$ where they increase by 2. They increase by 1 at the vertical arrows. Hence i commutative squares of this diagram lead to the canonical decomposition of the morphism $\sigma \circ \delta_i^\varepsilon$:

$$\begin{array}{ccccccc} \mathbb{I}^{k-1} & \xrightarrow{\delta_1^{\varepsilon 1}} & \mathbb{I}^k & \xrightarrow{\delta_2^{\varepsilon 2}} & \dots & \xrightarrow{\delta_{j_1-1}^{\varepsilon j_1-1}} & \mathbb{I}^{k+j_1-2} \xrightarrow{\delta_{j_1+1}^{\varepsilon j_1+1}} \dots \\ & & & & & \delta_{j_i-1}^{\varepsilon j_i-1} \downarrow & \\ & & \dots & \xrightarrow{\delta_{j_i-1}^{\varepsilon j_i-1}} & \mathbb{I}^{k+j_i-1} & \xrightarrow{\delta_{j_i}^{\varepsilon}} & \mathbb{I}^{k+j_i} \xrightarrow{\delta_{j_i+1}^{\varepsilon j_i+1}} \mathbb{I}^{k+j_i+1} \longrightarrow \dots \end{array}$$

The map σ at variables $(x_1, \dots, x_k)$, as a composition of maps $\delta_1^{\varepsilon 1}, \delta_2^{\varepsilon 2}, \dots, \delta_{n-k}^{\varepsilon n-k}$, has values

$$\sigma(x_1, \dots, x_k) = (\varepsilon_1, \dots, \varepsilon_{j_1-1}, x_1, \varepsilon_{j_1+1}, \dots, \varepsilon_{j_k-1}, x_k, \varepsilon_{j_k+1}, \dots, \varepsilon_{n-k}).$$

Its image is equals to $I_1 \times I_2 \times \dots \times I_n$ where $I_{j_1}, I_{j_2}, \dots, I_{j_k}$ are nondegenerated intervals. The image of the map $\sigma \circ \delta_i^\varepsilon$ is equal to

$$I_1 \times \dots \times I_{j_i-1} \times \{\varepsilon\} \times I_{j_i+1} \times \dots \times I_n.$$

The image $I_1 \times \dots \times I_n$ defines the map σ . Thus the differentials of the precubical set correspond to the differential of the chain complex $C_*(\mathcal{K}([0, 1]^m))$. Using Lemma 1.1 we obtain the required assertion. $\square$

REMARK 1.9 *Similar assertion remains true if we substitute $\square_+$ by a category Δ of finite totally ordered sets $[n] = \{0, 1, \dots, n\}$, $n \geq 0$ and nondecreasing maps. Differentials $d_k : L\Delta([k], [n]) \rightarrow L\Delta([k-1], [n])$ are defined as $d_k(\sigma) = \sum_{i=0}^k (-1)^i L\Delta(\delta_i^k, [n])(\sigma)$ where $\delta_i^k : [k-1] \rightarrow [k]$ is increasing maps such that $i \notin \text{Im}(\delta_i^k)$. Similar assertion also is true for a category Δ_+ of finite totally ordered sets $[n]$, $n \geq 0$, and increasing maps [18, Lemma 1.2]. But it is false for a category $\square$ with objects $\text{Ob}(\square) = \text{Ob}(\square_+)$ and morphisms generated by $\delta_i^{n,\varepsilon}$ and nondecreasing surjections $\sigma_i^n : \mathbb{I}^n \rightarrow \mathbb{I}^{n-1}$, $\sigma_i^n(x_1, \dots, x_n) = (x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n)$. Take for example $n = 0$. Then we obtain a chain complex*

$$0 \leftarrow \mathbb{Z} \xleftarrow{0} \mathbb{Z} \xleftarrow{0} \mathbb{Z} \xleftarrow{0} \dots$$

which all homology groups are isomorphic to $\mathbb{Z}$.

A prototype of the following proposition is [5, Lemma 4.2] which is proved there for simplicial objects of an abelian category. We prove it for precubical abelian groups.

Proposition 1.12 *Let $F : \square_+^{op} \rightarrow \text{Ab}$ be a precubical abelian group and F_* a chain complex $F_n = F(\mathbb{I}^n)$ with differentials $d_n = \sum_{i=1}^n (-1)^i (F(\delta_i^{n,1}) - F(\delta_i^{n,0}))$. Then for all $n \geq 0$ there are isomorphisms $\varinjlim_n^{\square_+^{op}} F \cong H_n(F_*)$ natural in $F \in \text{Ab}^{\square_+^{op}}$.*

PROOF. It follows from Lemma 1.11 that the sequence

$$0 \leftarrow \Delta_{\square_+} \mathbb{Z} \xleftarrow{d_0} Lh^{\mathbb{I}^0} \xleftarrow{d_1} Lh^{\mathbb{I}^1} \leftarrow \dots \xleftarrow{d_k} Lh^{\mathbb{I}^k} \xleftarrow{d_{k+1}} \dots$$

is the projective resolution of $\Delta_{\square_+} \mathbb{Z} \in \text{Ab}^{\square_+}$. We have isomorphisms by Lemma 1.6 $\varinjlim_n F \cong H_n(P_* \otimes F)$. It follows from Lemma 1.5 that there are natural isomorphisms $Lh^{\mathbb{I}^n} \otimes F \xrightarrow{\cong} F(\mathbb{I}^n)$. The differentials of the chain complex $Lh^{\mathbb{I}^*} \otimes F$ are equal to $\sum_{i=1}^n (-1)^i (Lh^{\delta_i^{n,1}} - Lh^{\delta_i^{n,0}}) \otimes F$. The isomorphism $Lh^{(-)} \otimes F \xrightarrow{\cong} F(-)$ carries these differentials to homomorphisms $\sum_{i=1}^n (-1)^i (F(\delta_i^{n,1}) - F(\delta_i^{n,0}))$. From this we obtain isomorphisms $\varinjlim_n^{\square_+^{op}} F \cong H_n(F_*)$ which are natural in F . $\square$

In accordance with Remark 1.9 this proposition is true in the cases of the categories Δ and Δ_+ . The category $\square$ has the terminal object $\mathbb{I}^0$. Hence the functor $\varinjlim^{\square}$ is exact and we obtain by Lemma 1.4 that $\varinjlim_n^{\square_+^{op}} \Delta \mathbb{Z} \cong \varinjlim_n^{\square} \Delta \mathbb{Z} = 0$ for $n > 0$. We saw that $H_n(\Delta \mathbb{Z}_*) \cong \mathbb{Z}$ for all $n \in \mathbb{N}$ in this case. Consequently this proposition is false for the category $\square$.

Homology groups with coefficients in a homological system. Now we study homology groups $\varinjlim_n^{(\square_+/X)^{op}} F$ of a precubical set $X \in \text{Set}^{\square_+^{op}}$ with coefficients in a homological system F on X . Consider abelian groups $C_n(X, F) = \bigoplus_{\sigma \in X_n} F(\sigma)$. Define differentials $d_i^{n,\varepsilon} : C_n(X, F) \rightarrow C_{n-1}(X, F)$ as homomorphisms making the following diagrams commutative

$$\begin{array}{ccc} \bigoplus_{\sigma \in X_n} F(\sigma) & \xrightarrow{d_i^{n,\varepsilon}} & \bigoplus_{\sigma \in X_{n-1}} F(\sigma) \\ \text{\scriptsize in_σ} \uparrow & & \uparrow \text{\scriptsize $in_{\sigma \delta_i^\varepsilon}$} \\ F(\sigma) & \xrightarrow{F(\delta_i^\varepsilon : \sigma \delta_i^\varepsilon \rightarrow \sigma)} & F(\sigma \delta_i^\varepsilon) \end{array}$$

DEFINITION 1.10 Let $F : (\square_+/X)^{op} \rightarrow \text{Ab}$ be a homological system of abelian groups on a precubical set X . Homology groups $H_n(X, F)$ with coefficients in F are n -th homology groups of the chain complex $C_*(X, F)$ consisting of abelian groups $C_n(X, F) = \bigoplus_{\sigma \in X_n} F(\sigma)$ and differentials $d_n =$

$$\sum_{i=1}^n (-1)^i (d_i^{n,1} - d_i^{n,0}).$$

EXAMPLE 1.11 Consider functors $\mathbb{Z}^0, \mathbb{Z}^1 : \square_+^{op} \rightarrow \text{Ab}$ with values $\mathbb{Z}^\alpha(\mathbb{I}^n) = \mathbb{Z}$ at objects for all $n \in \mathbb{N}$. Its values at morphisms are defined by homomorphisms $\mathbb{Z}^\alpha(\delta_i^{n,\varepsilon}) : \mathbb{Z} \rightarrow \mathbb{Z}$ which equal $1_{\mathbb{Z}}$ for $\varepsilon = \alpha$ and 0 for $\varepsilon \neq \alpha$. For any precubical set $X = (X_n, \partial_i^{n,\varepsilon})$ are defined homological systems $\mathbb{Z}^0 \circ Q_X^{op}, \mathbb{Z}^1 \circ Q_X^{op} : (\square_+/X)^{op} \rightarrow \text{Ab}$ where $Q_X : \square_+/X \rightarrow \square_+$ is the forgetful functor of a fibre. We get two chain complexes $C_*(X, \mathbb{Z}^0 \circ Q_X^{op})$ and $C_*(X, \mathbb{Z}^1 \circ Q_X^{op})$. Complexes $C_*(X, \mathbb{Z}^\alpha \circ Q_X^{op})$ for $\alpha \in \{0, 1\}$ consist of abelian groups

$$0 \leftarrow L(X_0) \xleftarrow{d_1^\alpha} L(X_1) \xleftarrow{d_2^\alpha} L(X_2) \leftarrow \dots$$

with differential $d_n^0 = - \sum_{i=1}^n (-1)^i L(\partial_i^{n,\varepsilon})$ and $d_n^1 = \sum_{i=1}^n (-1)^i L(\partial_i^{n,\varepsilon})$. Therefore every of two types homology groups considered by Goubault [8] are particular cases of the homology groups with coefficients in a homological system.

Let $\mathcal{HS}$ be a category each object of which is a pair (X, F) consisting of a precubical set X and homological system $F : (\square_+/X)^{op} \rightarrow \text{Ab}$. Morphisms $(X, F) \rightarrow (Y, G)$ are given by morphisms of precubical sets $f : X \rightarrow Y$ for which the following diagram is commutative

$$\begin{array}{ccc} (\square_+/X)^{op} & \xrightarrow{F} & \text{Ab} \\ & \searrow (\square_+/f)^{op} & \nearrow G \\ & (\square_+/Y)^{op} & \end{array}$$

Here $(\square_+/f)^{op}$ denotes a functor dual to $(\square_+/f)(\sigma) = f \circ \sigma$. Since the correspondence $(X, F) \mapsto \check{F}$ give the functor $(-): \mathcal{HS} \rightarrow \text{Ab}^{\square_+^{op}}$, we obtain functors $(X, F) \mapsto H_n(X, F)$ and $(X, F) \mapsto \varinjlim_n^{(\square_+/X)^{op}} F$.

Theorem 1.13 There are isomorphisms $H_n(X, F) \cong \varinjlim_n^{(\square_+/X)^{op}} F$ which are natural in $(X, F) \in \mathcal{HS}$ for all $n \geq 0$.

PROOF. We have $\varinjlim_n^{(\square_+/X)^{op}} F \cong \varinjlim_n^{\square^{op}} \check{F}$ by Proposition 1.10. It follows from the property of Kan extensions $\text{Lan}^S \circ \text{Lan}^T \cong \text{Lan}^{S \circ T}$ that this isomorphism is natural in $(X, F) \in \mathcal{HS}$. The chain complex $\check{F}_*$ is natural isomorphic to $C_*(X, F)$. It follows from Proposition 1.12 that $\varinjlim_n^{\square^{op}} \check{F} \cong H_n(\check{F}_*)$. The isomorphisms $\varinjlim_n^{\square^{op}} G \xrightarrow{\cong} H_n(G_*)$ are natural in precubical abelian groups $G \in \text{Ab}^{\square^{op}}$. Therefore we have the natural isomorphisms $\varinjlim_n^{\square^{op}} \check{F} \xrightarrow{\cong} H_n(\check{F}_*)$. Consequently there is a natural isomorphism $H_n(C_*(X, F)) \cong \varinjlim_n^{(\square_+/X)^{op}} F$. $\square$

EXAMPLE 1.12 Let $\square^0 = h_{\mathbb{I}^0} \in \text{Set}^{\square^{op}}$ be a cubical point. The category $\square/\square^0$ is isomorphic to $\square$. Hence $\varinjlim_n^{(\square/\square^0)^{op}} \Delta \mathbb{Z} \cong \varinjlim_n^{pt} \Delta \mathbb{Z}$. But $H_n(\square^0, \Delta \mathbb{Z}) \cong \mathbb{Z}$ for all $n \geq 0$. Consequently Theorem 1.13 is false for cubical sets.

A criterion of a homological isomorphism. Let $f : X \rightarrow Y$ be a morphism of precubical sets. We identify cubes $\sigma \in X_m$ with corresponding natural transformations $\tilde{\sigma} : h_{\mathbb{I}^m} \rightarrow X$ which are called by *singular cubes*. The functor $\square_+/f : \square_+/X \rightarrow \square_+/Y$ carries every cube $\sigma : h_{\mathbb{I}^m} \rightarrow X$ to the composition $f \circ \sigma$. It assigns to any morphism (δ, σ, τ) of the category $\square_+/X$ the morphism $(\delta, f \circ \sigma, f \circ \tau)$. Let $f^* : \text{Ab}^{(\square_+/Y)^{op}} \rightarrow \text{Ab}^{(\square_+/X)^{op}}$ be a functor assigning to each homological system F on Y the composition $F \circ (\square_+/f)^{op}$. The functor f^* acts at morphisms $\eta : F_1 \rightarrow F_2$ by Godement's law $f^*(\eta) = \eta * ((\square_+/f)^{op})$, so $f^*(\eta)_\sigma = \eta_{f \circ \sigma}$. It follows from Proposition 1.7 that there are canonical homomorphisms $\varinjlim_n^{(\square_+/X)^{op}} f^*(F) \rightarrow \varinjlim_n^{(\square_+/Y)^{op}} F$ which are natural in F . Thus we have the homomorphisms $H_n(X, f^*(F)) \rightarrow H_n(Y, F)$.

Let $\overleftarrow{f}(\sigma)$ be a precubical set defined by a pullback diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ f_\sigma \uparrow & & \uparrow \sigma \\ \overleftarrow{f}(\sigma) & \longrightarrow & h_{\mathbb{I}^n} \end{array} \quad (3)$$

The precubical set $\overleftarrow{f}(\sigma)$ is an *inverse image of the singular cube* σ . Since $(\square_+/f)/\sigma \cong \square_+/\overleftarrow{f}(\sigma)$ and $\varinjlim_n^{(\square_+/\overleftarrow{f}(\sigma))^{op}} \Delta \mathbb{Z} \cong H_n(\overleftarrow{f}(\sigma), \Delta \mathbb{Z})$, it follows from Lemma 1.4 and Proposition 1.7 the following

Corollary 1.14 *The homomorphisms $H_n(X, f^*(F)) \rightarrow H_n(Y, F)$ are isomorphisms at all $F \in \text{Ab}^{(\square_+/Y)^{op}}$ and $n \geq 0$ if and only if $H_n(\overleftarrow{f}(\sigma), \Delta \mathbb{Z})$ are isomorphic to $\varinjlim_n^{pt} \Delta \mathbb{Z}$ for all $\sigma \in \square_+/Y$ and $n \geq 0$.*

Homology groups of a colimit of precubical sets. Let $\mathcal{A}$ be an arbitrary category. Functors from small categories into $\mathcal{A}$ are called *diagrams of object of the category $\mathcal{A}$* . If a domain of a diagram is a small category J , then this diagram is denoted by $\{X^i\}_{i \in J}$ where X^i are its values at $i \in \text{Ob}(J)$. In particular, a *diagram of precubical sets* $\{X^i\}_{i \in J}$ consists of precubical sets X^i and morphisms $X^\alpha : X^i \rightarrow X^j$ which are given for $\alpha \in I(i, j)$ and satisfy to equations $X^{\alpha \circ \beta} = X^\alpha \circ X^\beta$ and $X^{1_i} = 1_{X^i}$.

Recall that a first quadrant spectral sequence $E_{p,q}^r$ [21] is said to *converge* to a graded abelian group $\{H_n\}$, and we write in this case $E_{p,q}^r \Rightarrow H_{p+q}$ if for each $n \in \mathbb{N}$ there is a sequence of subgroups

$$0 = F_{-1}H_n \subseteq F_0H_n \subseteq F_1H_n \subseteq \cdots \subseteq F_nH_n = H_n,$$

such that $F_pH_n/F_{p-1}H_n \cong E_{p,n-p}^\infty$ for all $0 \leq p \leq n$.

Proposition 1.15 *Let $\{X^i\}_{i \in J}$ be a diagram of precubical sets $X^i \in \text{Set}^{\square_+^{op}}$ such that*

$$\varinjlim_q^J \{L(X_n^i)\}_{i \in J} = 0, \text{ for any } n \in \mathbb{N} \text{ and } q > 0. \quad (4)$$

Let $\lambda_i : X^i \rightarrow \varinjlim^J \{X^i\}_{i \in J}$ is the colimit cone. Then for every homological system

$$F : (\square_+/\varinjlim^J \{X^i\}_{i \in J})^{op} \rightarrow \text{Ab}$$

there exists a first quadrant spectral sequence with $E_{p,q}^2 = \varinjlim_p^J \{H_q(X^i, \lambda_i^*(F))\}_{i \in J}$ which converges to graded abelian group $\{H_n(\varinjlim^J \{X^i\}_{i \in J}, F)\}$.

PROOF. Let $\mathcal{A}$ be an abelian category with exact products and $F : \square_+/\varinjlim^J \{X^i\}_{i \in J} \rightarrow \mathcal{A}$ a functor. Under the condition (4) by [16, Cor. 2.4] there is a spectral sequence

$$E_2^{p,q} = \lim_{J^{op}}^p \{\lim_{\square_+/X^i}^q F \circ (\square_+/\lambda_i)\}_{i \in J} \Rightarrow \lim_{\square_+/\varinjlim^J \{X^i\}_{i \in J}}^{p+q} F$$

Substitute $\mathcal{A} = \text{Ab}^{op}$. We have obtained a spectral sequence

$$E_{p,q}^2 = \varinjlim_p^J \{\varinjlim_q^{(\square_+/X^i)^{op}} \lambda_i^*(F)\}_{i \in J} \Rightarrow \{\varinjlim_{p+q}^{(\square_+/\varinjlim^J \{X^i\}_{i \in J})^{op}} F\}.$$

Theorem 1.13 leads us to the wanted spectral sequence. $\square$

DEFINITION 1.13 *Let X be a precubical set. A locally directed cover of X is a diagram of precubical sets $\{X^i\}_{i \in J}$ over a partially ordered set J satisfying the following three conditions*

- 1) *it assigns to each pair of members $i \leq j$ of the set J an inclusion $X^i \subseteq X^j$;*
- 2) *for each $n \in \mathbb{N}$ the equation $X_n = \bigcup_{i \in J} X_n^i$ holds true;*
- 3) *for each $\sigma \in X_n^i \cap X_n^j$ there exists $k \in J$ such that $k \leq i$ and $k \leq j$ and $\sigma \in X_n^k$.*

Corollary 1.16 *Let $\{X^i\}_{i \in J}$ be a locally directed cover of a precubical set X with inclusions λ_i inclusions $X^i \subseteq X$ and F a homological system of abelian groups on X . Then there exists a first quadrant spectral sequence with $E_{p,q}^2 = \varinjlim_p^J \{H_q(X^i, \lambda_i^*(F))\}_{i \in J}$ which converges to $\{H_n(X, F)\}$.*

PROOF. It follows from Theorem 1.13 and [16, Cor. 2.5]. $\square$

EXAMPLE 1.14 *Consider a cover of a cubical subset X of Euclidian space by the elementary cubes. These cubes put together a partially ordered aet J ordered by the inclusion. The diagram of these cubes and inclusions is the locally directed cover. For the homological system $F = \Delta \mathbb{Z}$ the spectral sequence collapses and gives isomorphisms $H_n(X, \Delta \mathbb{Z}) \cong \varinjlim_n^J \Delta \mathbb{Z} \quad n \in \mathbb{N}$.*

A spectral sequence of a morphism. Let $f : X \rightarrow Y$ be a morphism of cubical sets. For each $\sigma \in \square_+/Y$ the pullback diagram (3) defines the inverse image of the cube σ with the morphism $f_\sigma : \overleftarrow{f}(\sigma) \rightarrow X$. The inverse images of the singular cubes make up a diagram $\{\overleftarrow{f}(\sigma)\}_{\sigma \in \square_+/Y}$ with the colimit isomorphous to X . Using a spectral sequence of a morphism [17, 4.1] where we substitute $\mathcal{A}$ by the category Ab^{op} obtain the following assertion by Theorem 1.13:

Corollary 1.17 *Let $f : X \rightarrow Y$ be a morphism of precubical sets and $F : (\square_+/X)^{op} \rightarrow \text{Ab}$ homological system of abelian groups on X . Then there exists a first quadrant spectral sequence*

$$E_{p,q}^2 = \varinjlim_p^{\square_+/Y} \{H_q(\overleftarrow{f}(\sigma), f_\sigma^*(F))\}_{\sigma \in \square_+/Y} \Rightarrow H_{p+q}(X, F).$$

In particular, if the diagram $\{H_q(\overleftarrow{f}(\sigma), f_\sigma^*(F))\}_{\sigma \in \square_+/Y}$ consists of isomorphisms, then we can to obtain a diagram over the dual category by inverting

the isomorphisms. Denote it by $\{H_q(\overleftarrow{f}(\sigma), f_\sigma^*(F))\}_{\sigma \in (\square_+/Y)^{op}}^{-1}$. Using Theorem 1.13 we get the spectral sequence

$$E_{p,q}^2 \cong H_p(Y, \{H_q(\overleftarrow{f}(\sigma), f_\sigma^*(F))\}_{\sigma \in (\square_+/Y)^{op}}^{-1}) \Rightarrow H_{p+q}(X, F),$$

concerning homology of X and Y .

1.5 The category of factorizations

Let $\mathcal{C}$ be a small category. Denote by $\mathfrak{F}\mathcal{C}$ the *category of factorization in $\mathcal{C}$* [1]. Recall that $\text{Ob}(\mathfrak{F}\mathcal{C}) = \text{Mor}(\mathcal{C})$ and for every $\alpha, \beta \in \text{Ob}(\mathfrak{F}\mathcal{C})$ the set of morphisms $\mathfrak{F}\mathcal{C}(\alpha, \beta)$ consists of the pairs (f, g) of morphisms $f, g \in \text{Mor}\mathcal{C}$

$$\begin{array}{ccc} \circ & \xrightarrow{g} & \circ \\ \uparrow \alpha & & \uparrow \beta \\ \circ & \xleftarrow{f} & \circ \end{array}$$

satisfying $g \circ \alpha \circ f = \beta$. The composition of $\alpha \xrightarrow{(f_1, g_1)} \beta$ and $\beta \xrightarrow{(f_2, g_2)} \gamma$ is defined by $\alpha \xrightarrow{(f_1 \circ f_2, g_2 \circ g_1)} \gamma$. The identity morphism of an object $a \xrightarrow{\alpha} b$ in $\mathfrak{F}\mathcal{C}$ is the pair of the identity morphisms $\alpha \xrightarrow{(1_a, 1_b)} \alpha$.

Lemma 1.18 *For any category $\mathcal{C}$ there is an isomorphism $\mathfrak{F}(\mathcal{C}^{op}) \cong \mathfrak{F}\mathcal{C}$.*

Corollary 1.19 *Let $\mathcal{C}$ be a category and α its morphism. $\mathfrak{F}\mathcal{C}/\alpha$ is isomorphic to the category with objects triples of morphisms (x, β, y) for which the composition is defined and $x \circ \beta \circ y = \alpha$ and with morphisms $(x, \beta, y) \rightarrow (x', \beta', y')$ commutative diagrams*

$$\begin{array}{ccccc} & & \circ & \xleftarrow{\beta} & \circ \\ & x \swarrow & \downarrow v & & \nwarrow y \\ \circ & & & & \circ \\ & \swarrow x' & \downarrow w & & \searrow y' \\ & & \circ & \xleftarrow{\beta'} & \circ \end{array}$$

In [1] it was introduced the cohomology of categories with coefficients in natural system. It follows from [1, Theorem 4.4] that this cohomology may be defined as the right derived functor $\varprojlim_{\mathfrak{F}\mathcal{C}}^n$ of limit. We give a dual definition.

DEFINITION 1.15 *Let $\mathcal{C}$ be a small category and $\mathfrak{F}\mathcal{C}$ the category of factorizations. A contravariant natural system on $\mathcal{C}$ is a functor $F : (\mathfrak{F}\mathcal{C})^{op} \rightarrow \text{Ab}$. For $n \geq 0$ the n -th homology group of $\mathcal{C}$ with coefficients in F is the abelian groups $\varinjlim_n^{(\mathfrak{F}\mathcal{C})^{op}} F$.*


Since the category of factorization has a complicated structure, we seek its strong coinital subcategories with simple building. To this end we will study comma categories $\mathfrak{F}\mathcal{C}/\alpha$.

A category is said to be *cancellative* if its morphisms are epimorphic and monomorphic. A category $\mathcal{C}$ is cancellative if and only if the implications $\alpha\gamma = \beta\gamma \Rightarrow \alpha = \beta$ and $\gamma\alpha = \gamma\beta \Rightarrow \alpha = \beta$ are true for all $\alpha, \beta, \gamma \in \mathcal{C}$. The following lemma is useful for cancellative categories.

Recall that a preordered set may be defined as a small category $\mathcal{C}$ such that for any $a, b \in \text{Ob } \mathcal{C}$ the set $\mathcal{C}(a, b)$ is either empty or has precisely one element. In the latter case we write $a \leq b$. If moreover $a \leq b$ and $b \leq a$ implies $a = b$, then $\mathcal{C}$ is the partially ordered set.

Lemma 1.20 *Let $\mathcal{C}$ be a small cancellative category. If $\mathcal{C}$ does not contain nonidentity retractions, then the comma category $\mathfrak{F}\mathcal{C}/\alpha$ is the partially ordered set for each $\alpha \in \text{Ob}(\mathfrak{F}\mathcal{C})$.*

PROOF. For any $\alpha \in \text{Mor } \mathcal{C}$ the objects of the category $\mathfrak{F}\mathcal{C}/\alpha$ are triples of morphisms (x, w, y) of $\mathcal{C}$ for which $y \circ w \circ x = \alpha$. Consider any (parallel) pair of morphisms $(x, w, y) \xrightarrow[(u_2, v_2)]{(u_1, v_1)} (x', w', y')$ of the category $\mathfrak{F}\mathcal{C}/\alpha$. These morphisms are given by the commutative diagram



It follows from $y'v_1 = y = y'v_2$ that $v_1 = v_2$. Similarly, $u_1x' = x = u_2x'$ implies $u_1 = u_2$. Hence $u_1 = u_2$ and $v_1 = v_2$. Consequently, any two morphisms $(x, w, y) \rightarrow (x', w', y')$ are same. It follows that $\mathfrak{F}\mathcal{C}/\alpha$ is the preordered set. If $\mathcal{C}$ does not contain nonidentity retraction, then any two

morphisms

$$(x, w, y) \begin{matrix} \xrightarrow{(u_1, v_1)} \\ \xleftarrow{(u_2, v_2)} \end{matrix} (x', w', y')$$

lead to the equalities $u_1x' = x$, $u_2x = x'$, $y'v_1 = y$, $yv_2 = y'$. It follows that $u_1u_2x = x$, $yv_2v_1 = y$. Since $\mathcal{C}$ is cancellative, we obtain that u_1u_2 and v_2v_1 are identity morphisms. If $\mathcal{C}$ does not contain nonidentity retraction, then it results from this that u_1 , u_2 , v_1 , v_2 are identity morphisms. Therefore the relation $\leq$ is antisymmetric and $\mathfrak{F}\mathcal{C}/\alpha$ is the partially ordered set. $\square$

1.6 Acyclic partially ordered sets

Recall that a small category $\mathcal{C}$ is *acyclic* if $H_n(\mathcal{C}) \cong H_n(pt)$ for all $n \geq 0$. The study of strong coinital functors is closely related with the acyclic property. We point out two simple conditions of the acyclic property of a partially ordered set.

Lemma 1.21 *Let X be a partially ordered set. Let V and W be its closed subsets for which $X = V \cup W$. If V , W , and $V \cap W$ are acyclic, then X is acyclic.*

PROOF. Let $C_*(X) = C_*(X, \Delta_X \mathbb{Z})$. The unique functor $X \rightarrow pt$ gives the chain homomorphism $C_*(X) \rightarrow C_*(pt)$. Denote by $\tilde{C}_*(X)$ its kernel. It is easy to see that X is acyclic if and only if $H_n(\tilde{C}_*(X)) = 0$ for all $q \geq 0$. The subsets V^{op} and W^{op} are open and $X^{op} = V^{op} \cup W^{op}$. There exists a exact sequence

$$0 \rightarrow \tilde{C}_*(V^{op} \cap W^{op}) \rightarrow \tilde{C}_*(V^{op}) \oplus \tilde{C}_*(W^{op}) \rightarrow \tilde{C}_*(X^{op}) \rightarrow 0.$$

A corresponding long exact sequence leads us to isomorphisms $H_n(\tilde{C}_*(X^{op})) = 0$ for all $q \geq 0$. $\square$

An easy induction proves

Corollary 1.22 *Let $X = \bigcup_{i=1}^n W_i$ be the union of closed subsets $W_i \subseteq X$. If the intersections $W_{i_1} \cap W_{i_2} \cap \cdots \cap W_{i_k}$ are acyclic for all $\{i_1, i_2, \dots, i_k\} \subseteq \{1, 2, \dots, n\}$, then X is acyclic.*

2 Homology groups of free partially commutative monoids

From here n -th integral homology groups $\varinjlim_n^{\mathcal{C}} \Delta \mathbb{Z}$ of a small category $\mathcal{C}$ will be denoted by $H_n(\mathcal{C})$. At first we consider the case of commutative monoids and then generalize obtained assertions to partially commutative monoids.

2.1 Homology of a commutative monoid

Factorization category of the free commutative monoid.

DEFINITION 2.1 *A n -th Leech homology group $H_n(M, F)$ with coefficients in a contravariant natural system $F : (\mathfrak{F}M)^{op} \rightarrow \text{Ab}$ is a group $\varinjlim_n^{(\mathfrak{F}M)^{op}} F$.*

Lemma 2.1 *Let $\mathbb{N} = \{1, a, a^2, \dots\}$ be a free monoid with one generator a and $T = \{1, a\}$ its subset and $\mathfrak{F}T \subset \mathfrak{F}\mathbb{N}$ a full subcategory of the factorization category of M with the set of objects T . Then the inclusion $\mathfrak{F}T \subset \mathfrak{F}\mathbb{N}$ is strong coinital.*

PROOF. The category $\mathfrak{F}T$ consists of two objects 1 and a and two morphisms $1 \xrightarrow{(1,a)} a$ and $1 \xrightarrow{(a,1)} a$ except the identities. For any $p \in \mathbb{N}$ objects of $\mathfrak{F}T/a^p$ may be partitioned of the following two classes

- $1 \xrightarrow{(a^s, a^t)} a^p$ where $s + t = p$, for $p \geq 0$,
- $a \xrightarrow{(a^s, a^t)} a^p$ where $s + t = p - 1$, for $p \geq 1$.

The fibre $\mathfrak{F}T/a^p$ does not contain nonidentity morphisms between objects of the same classes. Between any two objects there exists one morphism at the most. Every nonidentity morphism of the fibre acts from an object of the first class to an object of the second class and decreases a degree by 1 either a first or second morphism. Therefore the category $\mathfrak{F}T/a^p$ is isomorphic to a category consisting of the morphisms

$$(1, a^p) \rightarrow (1, a^{p-1}) \leftarrow (a^1, a^{p-1}) \rightarrow \dots$$

$$\dots \leftarrow (a^s, a^t) \rightarrow (a^s, a^{t-1}) \leftarrow (a^{s+1}, a^{t-1}) \rightarrow \dots \leftarrow (a^p, 1)$$

It is well known (for example, see [18, Prop. 2.2] for the dual assertion) that homology groups of any partially ordered set $(P, \leq)$ with coefficients in

a functor $G : P \rightarrow \text{Ab}$ are isomorphic to homology groups of a subcomplex $C_*^+(P, G)$ of $C_*(P, G)$ (see Definition 1.5) which consists of abelian groups $C_n^+(P, G) = \bigoplus_{p_0 < \dots < p_n} G(p_0)$ where $p_0, \dots, p_n$ are members of P for which the strong inequalities $p_0 < \dots < p_n$ hold. If $G = \Delta \mathbb{Z}$ then the subcomplex consists of free abelian groups $\bigoplus_{p_0 < \dots < p_n} \mathbb{Z}$ and differentials defined on generators by $d_n(p_0 < \dots < p_n) = \sum_{i=0}^n (-1)^i (p_0 < \dots < \widehat{p_i} < \dots < p_n)$. From here one can obtain the following chain complex for the calculating the integral homology groups of the partial set $\mathfrak{F}T/a^p$

$$0 \rightarrow \mathbb{Z}^{2p} \xrightarrow{d_1} \mathbb{Z}^{2p+1} \rightarrow 0,$$

The differential d_1 has the matrix

$$D_1 = \begin{pmatrix} -1 & 0 & 0 & \dots & 0 & 0 \\ 1 & 1 & 0 & \dots & 0 & 0 \\ 0 & -1 & -1 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 1 & 1 \\ 0 & 0 & 0 & \dots & 0 & -1 \end{pmatrix}$$

consisting of $2p+1$ strings and $2p$ columns. Since for every column $x \in \mathbb{Z}^{2p}$ the equation $D_1 x = 0$ has the unique solution $x = 0$, the homomorphism d_1 is injective. Hence the homology groups equal $H_1(\mathfrak{F}T/a^p) = 0$, $H_0(\mathfrak{F}T/a^p) = \mathbb{Z}$. $\square$

Consider the free commutative monoid $\mathbb{N}^n$ generated by $a_1, a_2, \dots, a_n$. Let $T^n \subset \mathbb{N}^n$ be a subset consisting of finite productions $a_{i_1} a_{i_2} \dots a_{i_k}$ for which $1 \leq i_1 < i_2 < \dots < i_k \leq n$. The category $\mathfrak{F}\mathbb{N}^n$ contains a full subcategory with a set of objects T^n . Denote this subcategory by $\mathfrak{F}T^n$.

Lemma 2.2 *The inclusion $\mathfrak{F}T^n \subset \mathfrak{F}\mathbb{N}^n$ is strong coinital.*

PROOF. For every $\alpha = (a_1^{p_1}, a_2^{p_2}, \dots, a_n^{p_n})$ objects of the category $\mathfrak{F}T^n/\alpha$ may be considered as morphisms

$$(a_1^{\varepsilon_1}, a_2^{\varepsilon_2}, \dots, a_n^{\varepsilon_n}) \rightarrow (a_1^{p_1}, a_2^{p_2}, \dots, a_n^{p_n}),$$

where $\varepsilon_i \in \{0, 1\}$. The following commutative triangles correspond to it morphisms

$$\begin{array}{ccc} (a_1^{\varepsilon_1}, a_2^{\varepsilon_2}, \dots, a_n^{\varepsilon_n}) & \xrightarrow{\quad\quad\quad} & (a_1^{p_1}, a_2^{p_2}, \dots, a_n^{p_n}) \\ & \searrow \quad \quad \nearrow & \\ & (a_1^{\varepsilon'_1}, a_2^{\varepsilon'_2}, \dots, a_n^{\varepsilon'_n}) & \end{array}$$

where $\varepsilon_i \in \{0, 1\}$, $\varepsilon'_i \in \{0, 1\}$, for $1 \leq i \leq n$. We conclude from this that $\mathfrak{F}T^n/\alpha \cong (\mathfrak{F}T_1/a_1^{p_1}) \times \dots \times (\mathfrak{F}T_n/a_n^{p_n})$ where $\mathfrak{F}T_i$ a category consisting of two the objects 1 and a_i and two the morphisms $1 \xrightarrow{(1, a_i)} a_i$ and $1 \xrightarrow{(a_i, 1)} a_i$ except the identity morphisms. Using Lemma 2.1 we get the isomorphisms $H_q(\mathfrak{F}T^n/\alpha) \cong H_q(pt)$. Consequently, the inclusion $\mathfrak{F}(T^n) \rightarrow \mathfrak{F}\mathbb{N}^n$ is strong coinitial. $\square$

Leech homology as a cubical homology. Consider a precubical set T_*^n consisting of sets

$$T_k^n = \{a_{i_1}a_{i_2} \cdots a_{i_k} : 1 \leq i_1 < i_2 < \cdots < i_k \leq n\}$$

and maps, for $1 \leq s \leq k$,

$$T_{k-1}^n \xrightleftharpoons[\partial_s^1]{\partial_s^0} T_k^n, \quad \partial_s^0(a_{i_1} \cdots a_{i_k}) = \partial_s^1(a_{i_1} \cdots a_{i_k}) = a_{i_1} \cdots a_{i_{s-1}} \widehat{a_{i_s}} a_{i_{s+1}} \cdots a_{i_k}$$

Objects of the category $\square_+/T_*^n$ may be considered as pairs (k, a) where $a \in T_k^n$. Each $a \in T_k^n$ has an unique decomposition $a_{i_1} \cdots a_{i_k}$ for which $1 \leq i_1 < \cdots < i_k \leq n$. Thus the objects (k, a) may be identified with the elements $a_{i_1} \cdots a_{i_k} \in T^n$. Morphisms of $\square_+/T_*^n$ are triples $(\delta : I^m \rightarrow I^k, a_{j_1} \cdots a_{j_m}, a_{i_1} \cdots a_{i_k})$ satisfying $T^n(\delta)(a_{j_1} \cdots a_{j_m}) = a_{i_1} \cdots a_{i_k}$.

It allows us to build a functor $\mathfrak{S} : \square_+/T_*^n \rightarrow \mathfrak{F}T^n$ as follows. Define at objects $\mathfrak{S}(a_{i_1} \cdots a_{i_k}) = a_{i_1} \cdots a_{i_k}$. Define $\mathfrak{S}$ at morphisms as

$$\mathfrak{S}(\delta_s^{k,0}, a_{i_1} \cdots \widehat{a_{i_s}} \cdots a_{i_k}, a_{i_1} \cdots a_{i_k}) = (a_{i_s}, 1) : a_{i_1} \cdots \widehat{a_{i_s}} \cdots a_{i_k} \rightarrow a_{i_1} \cdots a_{i_k},$$

$$\mathfrak{S}(\delta_s^{k,1}, a_{i_1} \cdots \widehat{a_{i_s}} \cdots a_{i_k}, a_{i_1} \cdots a_{i_k}) = (1, a_{i_s}) : a_{i_1} \cdots \widehat{a_{i_s}} \cdots a_{i_k} \rightarrow a_{i_1} \cdots a_{i_k}.$$

Every morphism of the category $\square_+/T_*^n$ has a decomposition of the form $(\delta_s^{k,\varepsilon}, a_{i_1} \cdots \widehat{a_{i_s}} \cdots a_{i_k}, a_{i_1} \cdots a_{i_k})$, $\varepsilon \in \{0, 1\}$. There are the commutative dia-

grams for all $1 \leq s < t \leq k$ and $\alpha, \beta \in \{0, 1\}$

$$\begin{array}{ccc}
a_{i_1} \cdots \widehat{a_{i_s}} \cdots \widehat{a_{i_t}} \cdots a_{i_k} & \xrightarrow{\mathfrak{S}(\delta_s^\alpha)} & a_{i_1} \cdots \widehat{a_{i_t}} \cdots a_{i_k} \\
\mathfrak{S}(\delta_{t-1}^\beta) \downarrow & & \downarrow \mathfrak{S}(\delta_t^\beta) \\
a_{i_1} \cdots \widehat{a_{i_s}} \cdots a_{i_k} & \xrightarrow{\mathfrak{S}(\delta_s^\alpha)} & a_{i_1} \cdots a_{i_k}
\end{array}$$

It follows from this that the map $\mathfrak{S}$ defined above at morphisms as

$$(\delta_s^{k,\varepsilon}, a_{i_1} \cdots \widehat{a_{i_s}} \cdots a_{i_k}, a_{i_1} \cdots a_{i_k}),$$

has a unique extension by a functor. Here $\mathfrak{S}(\delta)$ is a short denotation of the values of $\mathfrak{S}$ at the morphism $(\delta : I^m \rightarrow I^k, a_{j_1} \cdots a_{j_m}, a_{i_1} \cdots a_{i_k})$. It is easy to see that the values $\mathfrak{S}(\delta)$ consist of pairs of morphisms (f, g) where f is the value of $\mathfrak{S}$ at the composition of morphisms of the form δ_s^0 appeared in the canonical decomposition of δ and g is the similar value at the composition of δ_s^1 .

Lemma 2.3 *The functor $\mathfrak{S} : \square_+ / T_*^n \rightarrow \mathfrak{F}T^n$ is strong coinital.*

PROOF. Consider the category $\mathfrak{S}/a$. Every morphism $\mathfrak{S}(I^m) \rightarrow a$ has the unique decomposition

$$\begin{array}{ccc}
\mathfrak{S}(I^m) & \xrightarrow{(f,g)} & a \\
& \searrow \mathfrak{S}(\gamma) & \nearrow (1,1) \\
& \mathfrak{S}(I^k) &
\end{array}$$

Consequently $\mathfrak{S}/a \cong \square_+ / I^k$. The category $\square_+ / I^k$ has the terminal object. Hence it is connected and acyclic. Thus the functor $\mathfrak{S}$ is strong coinital. $\square$

We assign to an arbitrary contravariant system $F : (\mathfrak{F}\mathbb{N}^n)^{op} \rightarrow \text{Ab}$ a homological system $\overline{F} = F|_{(\mathfrak{F}T^n)^{op}} \circ \mathfrak{S}^{op}$ on the precubical set T_*^n .

Theorem 2.4 *Let $F : (\mathfrak{F}\mathbb{N}^n)^{op} \rightarrow \text{Ab}$ be a contravariant natural system of abelian groups on the monoid $\mathbb{N}^n$. Then for all $k \geq 0$ it holds isomorphisms of Leech homology groups and cubical homology groups*

$$H_k(\mathbb{N}^n, F) \cong H_k(T_*^n, \overline{F}).$$

PROOF. The inclusion $\mathfrak{F}T^n \subset \mathfrak{F}\mathbb{N}^n$ is strong cointial by Lemma 2.2 and hence $\varinjlim_n^{(\mathfrak{F}\mathbb{N}^n)^{op}} F \cong \varinjlim_n F|_{(\mathfrak{F}T^n)^{op}}$.

The functor $\mathfrak{S}$ is strong cointial by Lemma 2.3. We have got an isomorphism $\varinjlim_k^{(\mathfrak{F}\mathbb{N}^n)^{op}} F \cong \varinjlim_k^{(\square_+/T_*^n)^{op}} F|_{(\mathfrak{F}T^n)^{op}} \circ \mathfrak{S}^{op}$. The definition of the Leech homology groups with the cubical homology groups and Theorem 1.13 leads us to the wanted assertion. $\square$

Corollary 2.5 *Under the conditions of Theorem 2.4 Leech homology groups $H_k(\mathbb{N}^n, F)$ are isomorphic to the k -th homology groups of the chain complex*

$$0 \leftarrow F(1) \xleftarrow{d_1} \bigoplus_{1 \leq i_1 \leq n} F(a_{i_1}) \xleftarrow{d_2} \bigoplus_{1 \leq i_1 < i_2 \leq n} F(a_{i_1} a_{i_2}) \leftarrow \cdots \xleftarrow{d_n} F(a_1 a_2 \cdots a_n) \leftarrow 0,$$

where $a_1, a_2, \dots, a_n$ are generators of the monoid $\mathbb{N}^n$ and

$$d_k((i_1, \dots, i_k), \varphi) = \sum_{s=1}^k (-1)^s ((i_1, \dots, \widehat{i_s}, \dots, i_k), \varphi),$$

$$F(a_{i_1} \cdots \widehat{a_{i_s}} \cdots a_{i_k}) \xrightarrow{(1, a_{i_s})} a_{i_1} \cdots a_{i_k} \varphi - F(a_{i_1} \cdots \widehat{a_{i_s}} \cdots a_{i_k}) \xrightarrow{(a_{i_s}, 1)} a_{i_1} \cdots a_{i_k} \varphi$$

Corollary 2.6 *If for any $1 \leq i \leq n$ and $a = a_{i_1} \cdots a_{i_k}$, $1 \leq i_1 < i_2 < \cdots < i_k \leq n$ it is true an equality of homomorphisms*

$$F((1, a_i) : a \rightarrow a_i a) = F((a_i, 1) : a \rightarrow a_i a)$$

$$\text{for all } i \notin \{i_1, \dots, i_k\}, \text{ then } H_k(\mathbb{N}^n, F) = \bigoplus_{1 \leq i_1 < i_2 < \cdots < i_k \leq n} F(a_{i_1} a_{i_2} \cdots a_{i_k})$$

PROOF. We have $d_k = 0$ for all $1 \leq k \leq n$. This leads to required isomorphisms. $\square$

2.2 Leech dimension of a free partially commutative monoid

Now we compute a homological Leech dimension of a free partially commutative monoid.

Homological dimension of small categories. Let $\mathcal{C}$ be a small category and $\varinjlim_n^{\mathcal{C}} : \text{Ab}^{\mathcal{C}} \rightarrow \text{Ab}$ the left satellites of the colimit. A *homological dimension* of $\mathcal{C}$ is defined by

$$\text{hd } \mathcal{C} = \sup\{n \in \mathbb{N} : \varinjlim_n^{\mathcal{C}} \neq 0\}.$$

Here $\sup$ is taken in the totally ordered set $\{-1\} \cup \mathbb{N} \cup \{-\infty\}$. For example, $\text{hd } \mathcal{C} = -1$ if and only if $\mathcal{C} = \emptyset$.

An *open subcategory* of the category $\mathcal{C}$ is a full subcategory $\mathcal{D} \subseteq \mathcal{C}$ which contains with each its object $d \in \mathcal{D}$ all the objects $c \in \mathcal{C}$ for which there are morphisms $d \rightarrow c$. A subcategory $\mathcal{D} \subseteq \mathcal{C}$ is *closed* if $\mathcal{D}^{op}$ is open subcategory of $\mathcal{C}^{op}$.

Lemma 2.7 *Let $\mathcal{C} = \bigcup_{j \in J} \mathcal{C}_j$ is an union of some open subcategories of $\mathcal{C}$. Then $\text{hd } \mathcal{C} = \sup_{j \in J} \{\text{hd } \mathcal{C}_j\}$.*

PROOF. For any $a \in \mathcal{C}$ denote by $\mathcal{C}^a \subseteq \mathcal{C}$ a full subcategory of $\mathcal{C}$ which objects $b \in \mathcal{C}^a$ have morphisms $a \rightarrow b$. It is the smallest open subcategory containing a . It follows from Mitchell assertion [22, Corollary 10] that $\text{hd } \mathcal{C} = \sup_{a \in \text{Ob } \mathcal{C}} \{\text{hd } \mathcal{C}^a\}$. Since the subcategories $\mathcal{C}_j$ are open, it is true that $\mathcal{C}^a = \mathcal{C}_j^a$ for all $a \in \mathcal{C}_j$. The required equality follows from

$$\sup_j \left\{ \sup_{a \in \text{Ob } \mathcal{C}_j} \{\text{hd } \mathcal{C}^a\} \right\} = \sup_{a \in \bigcup_{j \in J} \text{Ob } \mathcal{C}_j} \{\text{hd } \mathcal{C}^a\}.$$

□

Category of factorizations of a free partially commutative monoid.

Let $M(E, I)$ be a free partially commutative monoid generated by a set E and relations $ab = ba$ which hold true for all $(a, b) \in I$. We say that a subset $S \subseteq E$ consists of *mutually commuting generators* if $(a, b) \in I$ for all $a, b \in S$.

Denote by $E_v \subseteq E$, $v \in V$, maximal subsets of the mutually commuting generators. Let $M(E_v) \subseteq M(E, I)$ be submonoids generated by E_v . Denote by $E_v \subseteq E$, $v \in V$, the maximal subsets consisting of its mutually independent generators. Let $M(E_v) \subseteq M(E, I)$ be denote the monoid generated by E_v .


The union $\bigcup_{v \in V} M(E_v) \subseteq M(E, I)$ is not a subcategory. It makes difficulties for the studying homology of $M(E, I)$. The idea to considering the inclusion of the corresponding categories of factorizations saves the situation: The full subcategory of $\mathfrak{F}M(E, I)$ with the set of objects $M(E_v)$ equals $\mathfrak{F}M(E_v)$. Moreover, the subcategories $\mathfrak{F}M(E_v) \subseteq \mathfrak{F}M(E, I)$ are closed.

First we study the comma category $\mathfrak{F}M(E, I)/\alpha$ for any element $\alpha \in M(E, I)$.

Proposition 2.8 *Let $M(E, I)$ be a free partially commutative monoid and $\alpha \in M(E, I)$ an element. Then the category $\mathfrak{F}M(E, I)/\alpha$ is a partially ordered set.*

PROOF. By [4, Corollary 2] if E is finite, then the monoid $M(E, I)$ is cancellative. Let E be infinite. Suppose that $yw_1x = yw_2x$. There is a finite subset $E' \subseteq E$ consisting of symbols which are multipliers of the elements $x, y, w_1, w_2 \in M(E, I)$. Since E' is finite, the submonoid $M(E', I \cap (E' \times E'))$ is cancellative. It follows that $w_1 = w_2$. Hence, $M(E, I)$ is cancellative. Assign to every $x \in M(E, I)$ a length $|x|$ of a word which presents x . It is called a *length* of x . It easy to see that $|xy| = |x| + |y|$ for all $x, y \in M(E, I)$. Therefore, $M(E, I)$ does not contain nonidentity retractions. By Lemma 1.20 we conclude that $\mathfrak{F}M(E, I)/\alpha$ is a partially ordered set. $\square$

EXAMPLE 2.2 Let $M(E, I)$ be a free partially commutative monoid and $a \in E$ a generator. Describe the partially ordered set $\mathfrak{F}M(E, I)/a$. It consists of the elements $(1, 1, a)$, $(1, a, 1)$, and $(a, 1, 1)$. Its morphisms are defined by the diagram



The morphisms $(1, 1, a) \xrightarrow{(1,a)} (1, a, 1)$ and $(a, 1, 1) \xrightarrow{(a,1)} (1, a, 1)$ leads us to the relation $(1, 1, a) < (1, a, 1) > (a, 1, 1)$ in the partially ordered set $\mathfrak{F}M(E, I)/a$.

Elements $x, y \in M(E, I)$ are said to be *commuting* if $xy = yx$. We notice attention that generators a and b are commuting if and only if $(a, b) \in I$ or $a = b$.

The category $\mathfrak{F}M(E, I)$ contains the full subcategory $\bigcup_{v \in V} \mathfrak{F}M(E_v)$ whose objects are products of the pairwise commuting generators.


Theorem 2.9 *The inclusion $\bigcup_{v \in V} \mathfrak{F}M(E_v) \subseteq \mathfrak{F}M(E, I)$ is strong coinitial.*

PROOF. By Lemma 1.18 the category $\mathfrak{F}M(E, I)$ is isomorphic to $\mathfrak{F}(M(E, I)^{op})$. This isomorphism carries the subcategory $\bigcup_{v \in V} \mathfrak{F}M(E_v)$ to $\bigcup_{v \in V} \mathfrak{F}(M(E_v)^{op})$.

Thus, it is enough to prove that the inclusion $\bigcup_{v \in V} \mathfrak{F}(M(E_v)^{op}) \subseteq \mathfrak{F}(M(E, I)^{op})$ is strong coinital. It is equivalent to the assertion that for each $\alpha \in M(E, I)$

$$H_q\left(\bigcup_{v \in V} \mathfrak{F}(M(E_v)^{op})/\alpha\right) \cong H_q(pt), \text{ for all } q \in \mathbb{N}.$$

We prove this assertion by induction on the length $|\alpha|$. By Proposition 2.8 the comma category $\mathfrak{F}(M(E, I)^{op})/\alpha$ is a partially ordered set. By Corollary 1.19 its elements may be given as triples (x, β, y) of elements of $M(E, I)$ satisfying $x \circ \beta \circ y = \alpha$. The relation $(x, \beta, y) \leq (x', \beta', y')$ is defined by existence of pairs (v, w) making the following diagram to be commutative



The subcategory $(\bigcup_{v \in V} \mathfrak{F}(M(E_v)^{op}))/\alpha \cong \bigcup_{v \in V} (\mathfrak{F}(M(E_v)^{op})/\alpha)$ is its closed subset consisting of the triples (x, β, y) for which $\beta \in \bigcup_{v \in V} M(E_v)$.

Let $\Phi(\alpha) = \bigcup_{v \in V} (\mathfrak{F}(M(E_v)^{op})/\alpha)$. We will prove by induction on $|\alpha|$ that the partially ordered set $\Phi(\alpha)$ is acyclic. We use that $\Phi(\alpha)$ is union of closed subsets which are isomorphic to $\Phi(\beta)$ with $|\beta| < |\alpha|$.

If $|\alpha| = 0$, then $\alpha = 1$, hence the partially ordered set $\Phi(\alpha)$ consists of unique element $(1, 1, 1)$. So, the assertion is true in this case.

Suppose that $\forall \beta \in M(E, I)$ with $|\beta| < n$ there are isomorphisms $H_q(\Phi(\beta)) \cong H_q(pt)$ for all $q \geq 0$. Prove it for α with $|\alpha| = n$.

If $\alpha = f \circ g$, then we say that f is a *left divisor* of $\alpha \in M(E, I)$ and g is its *right divisor*. In this case, there are injections of the partially ordered sets

$$\mathfrak{F}(M(E, I)^{op})/g \xrightarrow{f_*} \mathfrak{F}(M(E, I)^{op})/\alpha \xleftarrow{g^*} \mathfrak{F}(M(E, I)^{op})/f,$$

where $f_*(x, \beta, y) = (f \circ x, \beta, y)$, $g^*(x, \beta, y) = (x, \beta, y \circ g)$. Images of the maps

$$\Phi(g) \xrightarrow{f_*} \Phi(\alpha) \xleftarrow{g^*} \Phi(f)$$

are closed in $\Phi(\alpha)$. In general, if $f \circ \gamma \circ g = \alpha$ for some $f, g, \gamma \in M(E, I)$, then the image of the inclusion

$$f_* g^* = g^* f_* : \Phi(\gamma) \longrightarrow \Phi(\alpha)$$

is closed in $\Phi(\alpha)$.

Let $\{a_1, \dots, a_m\} \subseteq E$ be the set of generators which are left divisors of α and $\{b_1, \dots, b_n\} \subseteq E$ its right divisors. It is easy to see that the generators $a_1, \dots, a_m$ are mutually independent. Similarly $b_1, \dots, b_n$ are mutually independent. Denote by

$$L_i(\alpha) = \{(x, \beta, y) \in \Phi(\alpha) : x \text{ has the left divisor } a_i\}$$

$$R_j(\alpha) = \{(x, \beta, y) \in \Phi(\alpha) : y \text{ has the right divisor } b_j\}$$

for every $1 \leq i \leq m$ and $1 \leq j \leq n$. Let $L(\alpha) = \bigcup_{i=1}^m L_i(\alpha)$ and $R(\alpha) = \bigcup_{j=1}^n R_j(\alpha)$. It is evidently that $L(\alpha) = \{(x, \beta, y) : \beta \in \bigcup_{v \in V} M(E_v), x \neq 1\}$ and $R(\alpha) = \{(x, \beta, y) : \beta \in \bigcup_{v \in V} M(E_v), y \neq 1\}$.

Now consider the following two cases: $\alpha \in \bigcup_{v \in V} M(E_v)$ and $\alpha \notin \bigcup_{v \in V} M(E_v)$. If $\alpha \in \bigcup_{v \in V} M(E_v)$, then $\Phi(\alpha)$ contains the largest element $(1, \alpha, 1)$ and consequently $H_q(\Phi(\alpha)) \cong H_q(pt)$ for all $q \geq 0$.

If $\alpha \notin \bigcup_{v \in V} M(E_v)$, then $x \neq 1$ or $y \neq 1$ holds for each $(x, \beta, y) \in \Phi(\alpha)$. Therefore $(x, \beta, y) \in L(\alpha) \cup R(\alpha)$. It follows that $\Phi(\alpha) = L(\alpha) \cup R(\alpha)$.

Now we will prove the acyclic property of $L(\alpha)$, $R(\alpha)$, and $L(\alpha) \cap R(\alpha)$. Then, by Lemma 1.21, we will obtain the acyclic property of $L(\alpha) \cup R(\alpha)$. If $\alpha = f \circ g$, then denote g by $f^{-1}\alpha$ and f by αg^{-1} .

- $L(\alpha) = \bigcup_{i=1}^m L_i(\alpha)$, $L_{i_1}(\alpha) \cap \dots \cap L_{i_k}(\alpha) = (a_{i_1} \dots a_{i_k})_* \Phi((a_{i_1} \dots a_{i_k})^{-1}\alpha)$.

Since the map $(a_{i_1} \dots a_{i_k})_*$ makes an isomorphism of the partially ordered set $\Phi((a_{i_1} \dots a_{i_k})^{-1}\alpha)$ with a closed subset, the intersections $L_{i_1}(\alpha) \cap \dots \cap L_{i_k}(\alpha)$ are acyclic. By Corollary 1.22 we obtain the acyclic property of $L(\alpha)$.

- The acyclic property of $R(\alpha) = \bigcup_{j=1}^n R_j(\alpha)$ is proved similarly.

- $L(\alpha) \cap R(\alpha) = \bigcup_{i=1}^m (L_{i_1}(\alpha) \cap R(\alpha)), \quad L_{i_1}(\alpha) \cap \dots \cap L_{i_k}(\alpha) \cap R(\alpha) = \{(a_{i_1} \dots a_{i_k} x, \beta, y) : \beta \in \bigcup_{v \in V} M(E_v), y \neq 1, a_{i_1} \dots a_{i_k} x \beta y = \alpha\}$. It follows that

$$L_{i_1}(\alpha) \cap \dots \cap L_{i_k}(\alpha) \cap R(\alpha) = (a_{i_1} \dots a_{i_k})_* R((a_{i_1} \dots a_{i_k})^{-1} \alpha)$$

is isomorphic to the partially ordered set $R((a_{i_1} \dots a_{i_k})^{-1} \alpha)$ whose acyclic property is proved in the second item.

It follows from Lemma 1.21 that $\Phi(\alpha)$ is acyclic. $\square$

DEFINITION 2.3 *Let M be a monoid. Its a homological Leech dimension is defined by $\text{Ld } M = \text{hd } (\mathfrak{F}M)^{op}$.*

Corollary 2.10 *Let $M(E, I)$ be a free partially commutative monoid in which the maximal number of mutually commuting generators equals n . Then $\text{Ld } M(E, I) = n$.*

PROOF. Since $\bigcup_{v \in V} \mathfrak{F}M(E_v) \subseteq \mathfrak{F}M(E, I)$ is the closed subcategory, the following inequality follows from the assertion [22, Corollary 10]

$$\text{hd} \left(\bigcup_{v \in V} \mathfrak{F}M(E_v) \right)^{op} \leq \text{hd} (\mathfrak{F}M(E, I))^{op}.$$

The inclusion $\bigcup_{v \in V} \mathfrak{F}M(E_v) \subseteq \mathfrak{F}M(E, I)$ is strong coinital by Theorem 2.9.

Thus for any functor $F : (\mathfrak{F}M(E, I))^{op} \rightarrow \text{Ab}$ it follows from $\varinjlim_n^{(\mathfrak{F}M(E, I))^{op}} F \neq 0$ that $\varinjlim_n F|_{(\bigcup_{v \in V} \mathfrak{F}M(E_v))^{op}} \neq 0$. From here

$$\text{hd} \left(\bigcup_{v \in V} \mathfrak{F}M(E_v) \right)^{op} \geq \text{hd} (\mathfrak{F}M(E, I))^{op}.$$

Consequently, $\text{hd} \left(\bigcup_{v \in V} \mathfrak{F}M(E_v) \right)^{op} = \text{hd} (\mathfrak{F}M(E, I))^{op}$. From this, using Lemma 2.7 we get the equality $\text{hd} (\mathfrak{F}M(E, I))^{op} = \sup_{v \in V} \text{hd} (\mathfrak{F}M(E_v))^{op}$. So $\text{Ld } M(E, I) = \sup_{v \in V} \text{Ld } M(E_v)$. Since the homological Leech dimension of the free commutative monoid generated by n elements equals n , we conclude that $\text{Ld } M(E, I) = n$ where n is the maximal number of generators of $M(E_v)$. $\square$

2.3 Leech homology groups of free partially commutative monoids

A coinital subcategory of the factorization category. Let E be a set totally ordered by a relation $<$. We prove the following auxiliary assertion.

Proposition 2.11 *Let $T_v \subset M(E_v)$, for every $v \in V$, be a set consisting of products $a_1 a_2 \cdots a_n$ such that $a_1 < a_2 < \cdots < a_n$ and $a_j \in E_v$ for all $1 \leq j \leq n$. Here n is a finite number such that $n \leq |E_v|$. The product equals 1 $\in T_v$ for $n = 0$. Then the inclusion $\bigcup_{v \in V} \mathfrak{F}T_v \subset \mathfrak{F}M(E, I)$ is strong coinital.*

PROOF. A composition of strong coinital functors is strong coinital by Corollary 1.8. We have proved that the $\bigcup_{v \in V} \mathfrak{F}M(E_v) \subseteq \mathfrak{F}M(E, I)$ is strong coinital. Thus it is enough to show that the inclusion $\bigcup_{v \in V} \mathfrak{F}T_v \subset \bigcup_{v \in V} \mathfrak{F}M(E_v)$ is strong coinital. But for each $\alpha \in \bigcup_{v \in V} \mathfrak{F}M(E_v)$ there exists $w \in V$ such that $\alpha \in \mathfrak{F}M(E_w)$. All factors of α belong to $M(E_w)$. Consequently $\bigcup_{v \in V} \mathfrak{F}T_v / \alpha = \mathfrak{F}T_w / \alpha$.

The inclusion $\mathfrak{F}T_w \subset \mathfrak{F}M(E_w)$ is strong coinital by Lemma 2.2. So, $H_q(\bigcup_{v \in V} \mathfrak{F}T_v / \alpha) \cong H_q(pt)$. $\square$

Main Theorem. Let E be a set and $I \subseteq E \times E$ an irreflexive symmetric relation. We build a precubical set $T(E, I)$ depending on some total ordering relation $\leq$ on E . For every integer $n > 0$ define $T(E, I)_n$ as a set of words $a_1 \cdots a_n$ consisting of mutually commuting generators $a_1 < \cdots < a_n$ where $a_1, \dots, a_n \in E$

$$T(E, I)_n = \{a_1 \cdots a_n : (a_1 < \cdots < a_n) \& (1 \leq i < j \leq n \Rightarrow (a_i, a_j) \in I)\}.$$

(For $n = 0$ the set $T(E, I)_0$ consists of the empty word 1.) The maps $\partial_i^{n, \varepsilon} : T(E, I)_n \rightarrow T(E, I)_{n-1}$, for $1 \leq i \leq n$, act as

$$\partial_i^{n, 0}(a_1 \cdots a_n) = \partial_i^{n, 1}(a_1 \cdots a_n) = a_1 \cdots \widehat{a_i} \cdots a_n.$$

It easy to see that $T(E, I)$ is union of precubical sets $(T_v)_*$ for which

$$(T_v)_n = \{a_1 \cdots a_n \in T(E, I)_n : (1 \leq i \leq n \Rightarrow a_i \in E_v)\},$$

where $E_v \subseteq E$ are the maximal subsets of mutually commuting generators of the monoid $M(E, I)$. The maps $(T_v)_n \xrightarrow{\partial_i^{n, \varepsilon}} (T_v)_{n-1}$, act for $1 \leq i \leq n$ and $\varepsilon \in \{0, 1\}$ by

$$\partial_s^{n, 0}(a_1 \cdots a_n) = \partial_s^{n, 1}(a_1 \cdots a_n) = a_1 \cdots \widehat{a_s} \cdots a_n$$

Let $\mathfrak{S} : \square_+ / \bigcup_{v \in V} (T_v)_* \rightarrow \bigcup_{v \in V} \mathfrak{F}T_v$ be a functor which assign to every singular cube $a_1 \cdots a_n \in \bigcup_{v \in V} (T_v)_n$ the object $a_1 \cdots a_n$. The action of functor $\mathfrak{S}$ at morphisms is similar with the action in the case of free commutative monoid.

For any functor $F : (\mathfrak{F}M(E, I))^{op} \rightarrow \text{Ab}$ denote by $\overline{F}$ a homological system on $T(E, I)$ defined as the composition

$$(\square_+ / T(E, I))^{op} \xrightarrow{\mathfrak{S}^{op}} \bigcup_{v \in V} (\mathfrak{F}T_v)^{op} \subset (\mathfrak{F}M(E, I))^{op} \xrightarrow{F} \text{Ab}$$

DEFINITION 2.4 *A free partially commutative monoid $M(E, I)$ is called locally bounded if every set of mutually commuting generators is finite.*

Theorem 2.12 *Let $M(E, I)$ be a locally bounded free partially commutative monoid an $F : (\mathfrak{F}M(E, I))^{op} \rightarrow \text{Ab}$ a functor. Then for all $n \geq 0$ there are isomorphisms of the Leech and cubical homology groups*

$$H_n(M(E, I), F) \cong H_n(T(E, I), \overline{F}).$$

PROOF. The functor $\mathfrak{S}$ carries the subcategory $\square_+ / (T_v)_* \subseteq \square_+ / T(E, I)$ into $\mathfrak{F}T_v$ and hence defines a functor which we denote by $\mathfrak{S}_v : \square_+ / (T_v)_* \rightarrow \mathfrak{F}T_v$. For an arbitrary $\alpha \in \bigcup_{v \in V} \mathfrak{F}T_v$ there exists $w \in V$ such that $\alpha \in \mathfrak{F}T_w$.

Consequently there is an isomorphism $\mathfrak{S}/\alpha \cong \mathfrak{S}_w/\alpha$. It follows from Lemma 2.3 for the free finite generated commutative monoid $M(E_v)$ that the functor $\mathfrak{S}_v : \square_+ / (T_v)_* \rightarrow \mathfrak{F}T_v$ is strong coinital. So the category $\mathfrak{S}_v/\alpha$ is connected and acyclic. Therefore the category $\mathfrak{S}/\alpha$ is connected and acyclic. Thus $\mathfrak{S}$ is strong coinital. Using that the inclusion $\bigcup_{v \in V} \mathfrak{F}T_v \subseteq \mathfrak{F}M(E, I)$ is strong coinital we get a required isomorphisms

$$\varinjlim_n^{(\mathfrak{F}M(E, I))^{op}} F \cong \varinjlim_n^{\left(\square_+ / \bigcup_{v \in V} (T_v)_*\right)^{op}} \left(F|_{\bigcup_{v \in V} (\mathfrak{F}T_v)^{op}} \circ \mathfrak{S}^{op} \right)$$

□

From this we obtain the main result of the paper

Theorem 2.13 *Let $M(E, I)$ be a locally bounded free partially commutative monoid and $F : (\mathfrak{F}M(E, I))^{op} \rightarrow \text{Ab}$ a functor. Then groups $\varinjlim_n^{(\mathfrak{F}M(E, I))^{op}} F$ are isomorphic to n -th homology groups of a chain complex*

$$\begin{aligned} 0 \leftarrow F(1) \xleftarrow{d_1} \bigoplus_{a_1 \in E} F(a_1) \xleftarrow{d_2} \bigoplus_{\substack{a_1 < a_2 \\ (a_1, a_2) \in I}} F(a_1 a_2) \leftarrow \dots \\ \dots \leftarrow \bigoplus_{\substack{a_1 < a_2 < \dots < a_{n-1} \\ (a_i, a_j) \in I}} F(a_1 a_2 \dots a_{n-1}) \xleftarrow{d_n} \bigoplus_{\substack{a_1 < a_2 < \dots < a_n \\ (a_i, a_j) \in I}} F(a_1 a_2 \dots a_n) \leftarrow \dots, \end{aligned}$$

n -th member of which for $n \geq 0$ equals a direct sum of abelian groups $F(a_1 \dots a_n)$ taken by all sequences $a_1 < a_2 < \dots < a_n$ of mutually commuting generators $a_1, a_2, \dots, a_n \in E$ with the differentials

$$d_n((a_1, \dots, a_n), \varphi) = \sum_{s=1}^n (-1)^s ((a_1, \dots, \widehat{a}_s, \dots, a_n), F(1, a_s)\varphi - F(a_s, 1)\varphi)$$

where $F(1, a_s)$ shortly denote values F at morphisms $(1, a_s) : a_1 \dots \widehat{a}_s \dots a_n \rightarrow a_1 \dots a_n$, and $F(a_s, 1)$ at $(a_s, 1) : a_1 \dots \widehat{a}_s \dots a_n \rightarrow a_1 \dots a_n$.

PROOF. It follows from Theorem 2.12 that $\varinjlim_n^{(\mathfrak{F}M(E, I))^{op}} F \cong \varinjlim_n^{(\square_+ / T(E, I))^{op}} \overline{F}$. Theorem 1.13 leads to the isomorphisms

$$\varinjlim_n^{\mathfrak{F}M(E, I)^{op}} F \cong H_n(C_*(T(E, I), \overline{F})).$$

□

2.4 Hochschild homology

Lemma 2.14 *Let $S : \mathcal{C} \rightarrow \mathcal{D}$ be a functor between small categories. Then there exists a isomorphism which is natural in $F \in \text{Ab}^{\mathcal{C}}$ and $G \in \text{Ab}^{\mathcal{D}^{op}}$*

$$\text{Lan}^S F \otimes G \cong F \otimes (G \circ S).$$

PROOF. Recall that $\text{Hom}(F, A) : \mathcal{C}^{op} \rightarrow \text{Ab}$ for an abelian group A denotes a functor having values $\text{Ab}(F(c), A)$ at objects $c \in \mathcal{C}$ at $\text{Ab}(F(\alpha), 1_A)$ at morphisms $\alpha \in \text{Mor}(\mathcal{C})$. Since Lan^S is left adjoint to $(-) \circ S$, there are isomorphisms which are natural in $F \in \text{Ab}^{\mathcal{C}}$, $G \in \text{Ab}^{\mathcal{D}^{op}}$, and $A \in \text{Ab}$

$$\text{Ab}^{\mathcal{D}}(\text{Lan}^S F, \text{Hom}(G, A)) \cong \text{Ab}^{\mathcal{C}}(F, \text{Hom}(G, A) \circ S) \cong \text{Ab}^{\mathcal{C}}(F, \text{Hom}(G \circ S^{op}, A)).$$

It follows from isomorphisms $\text{Ab}(T \otimes G, A) \cong \text{Ab}^{\mathcal{D}}(T, \text{Hom}(G, A))$ which are natural in $T \in \text{Ab}^{\mathcal{D}}$ that there exists the isomorphism

$$\text{Ab}(\text{Lan}^S F \otimes G, A) \xrightarrow{\cong} \text{Ab}(F \otimes (G \circ S^{op}), A)$$

which is natural in three arguments. Since the Yoneda inclusion $T \mapsto \text{Ab}(T, -)$ is full, there exists an isomorphism $F \otimes (G \circ S) \xrightarrow{\cong} \text{Lan}^S F \otimes G$. $\square$

Let $(s, t) : \mathfrak{F}\mathcal{C} \rightarrow \mathcal{C}^{op} \times \mathcal{C}$ be a functor assigning to every object $a \xrightarrow{\alpha} b$ the pair $(a, b) \in \text{Ob}(\mathcal{C}^{op} \times \mathcal{C})$ and to every morphism $\alpha \xrightarrow{(f, g)} \beta$ the pair $(f, g) \in \text{Mor}(\mathcal{C}^{op} \times \mathcal{C})$.

Let $\mathcal{C}$ be a small category. We define its *Hochschild–Mitchell homology groups with coefficients in a $\mathcal{C}^{op}$ -bimodule $B : (\mathcal{C}^{op} \times \mathcal{C})^{op} \rightarrow \text{Ab}$* as $\text{Tor}_n(L\mathcal{C}, B)$ where $L\mathcal{C} : \mathcal{C}^{op} \times \mathcal{C} \rightarrow \text{Ab}$ is the composition of functors $\mathcal{C}^{op} \times \mathcal{C} \xrightarrow{\mathcal{C}(-;=)} \text{Set} \xrightarrow{L} \text{Ab}$.

Lemma 2.15 $\text{Tor}_n(L\mathcal{C}, B) \cong \varinjlim_n^{\mathfrak{F}\mathcal{C}^{op}} B \circ (s, t)^{op}$

PROOF. Since connected components of the category $(s, t)/(a, b)$ have terminal objects by [18, Lemma 2.4], the application of $\text{Lan}^{(s, t)}$ to a projective resolution $P_* \rightarrow \Delta \mathbb{Z}$ gives a projective resolution of $\text{Lan}^{(s, t)} \Delta \mathbb{Z} = L\mathcal{C}$. From the isomorphisms $\text{Lan}^{(s, t)} P_n \otimes B \cong P_n \otimes B \circ (s, t)^{op}$ we obtain the isomorphisms

$$\text{Tor}_n(\text{Lan}^{(s, t)} \Delta \mathbb{Z}, B) \cong \text{Tor}_n(\Delta \mathbb{Z}, B \circ (s, t)^{op})$$

The required assertion follows from $\text{Tor}_n(\Delta \mathbb{Z}, B \circ (s, t)^{op}) \cong \varinjlim_n^{\mathfrak{F}\mathcal{C}^{op}} B \circ (s, t)^{op}$. $\square$

Let M be a monoid considered as a category with one object. Then its Hochschild–Mitchell homology groups are isomorphic to Hochschild homology groups $H_n(\mathbb{Z}[M], B)$ of the monoid ring $\mathbb{Z}[M]$ with coefficients in $\mathbb{Z}[M] \otimes \mathbb{Z}[M]^{op}$ -module B .

Corollary 2.16 *Let $M(E, I)$ be a locally bounded free partially commutative monoid and B be a $\mathbb{Z}[M(E, I)] \otimes \mathbb{Z}[M(E, I)]^{op}$ -module. Then the Hochschild homology groups $H_n(\mathbb{Z}[M(E, I)], B)$ are isomorphic to homology groups of the chain complex consisting of the abelian groups $C_0 = B$ and $C_n = \bigoplus_{(a_1, \dots, a_n)} B$ for $n > 0$ where the sums are taken by all sequences of mutually commuting generators $a_1 < \dots < a_n$, $a_1, \dots, a_n \in E$. Differentials*

are defined by the following values, for $\varphi \in B$,

$$d_n(a_1, \dots, a_n, \varphi) = \sum_{i=1}^n (-1)^i (a_1, \dots, \widehat{a_i}, \dots, a_n, a_i \cdot \varphi - \varphi \cdot a_i).$$

Here the dot denotes both a left and a right action of a monoid on B .

(These actions may be written as $a \cdot \varphi \cdot b = B(a, b)(\varphi)$ by the bifunctor $B(-, =)$.)

2.5 Homology with coefficients in a right module

Let $\mathcal{C}$ be an arbitrary small category. Consider a functor $\mathfrak{F}\mathcal{C} \xrightarrow{t} \mathcal{C}$ assigning to every arrow $a \rightarrow b$ the end b and to every morphism $\alpha \xrightarrow{(f, g)} \beta$ the morphism $g : t(\alpha) \rightarrow t(\beta)$. The following assertion is implicitly contained in [18].

Lemma 2.17 *The functor $\mathfrak{F}\mathcal{C} \xrightarrow{t} \mathcal{C}$ is strong coinital.*

PROOF. For each object $c \in \mathcal{C}$ the category t/c contains a full subcategory $\mathcal{D}$ consisting of all objects $(\beta, 1_c)$. This subcategory is isomorphic to $(\mathcal{C}/c)^{op}$. For any object (α, x) it is defined a morphism from (α, x) to $(x \circ \alpha, 1_c) \in Ob(\mathcal{D})$ given by the diagram

$$\begin{array}{ccc} a & \xrightarrow{\alpha} & b \\ \uparrow 1_a & & \downarrow x \\ a & \xrightarrow{x \circ \alpha} & c \end{array} \quad \begin{array}{c} \searrow x \\ \nearrow 1_c \end{array} \quad c$$

The morphism $(\alpha, x) \rightarrow (x \circ \alpha, 1_c)$ is universal in the sense that for each morphism $(\alpha, x) \rightarrow (\beta, 1_c)$ there exists an unique morphism $(x \circ \alpha, 1_c) \rightarrow (\beta, 1_c)$ making commutative the following diagram

$$\begin{array}{ccc} (\alpha, x) & \xrightarrow{\quad} & (x \circ \alpha, 1_c) \\ & \searrow & \swarrow \exists! \\ & (\beta, 1_c) & \end{array}$$

Consequently, there is a functor $\sigma : t/c \rightarrow (\mathcal{C}/c)^{op}$ which is left adjoint to a full inclusion $(\mathcal{C}/c)^{op} \subseteq t/c$. Hence the inclusion $\mathcal{C}/c \subseteq (t/c)^{op}$ is strong coinital. Therefore $\varinjlim_n^{t/c}(\Delta \mathbb{Z}) \circ \sigma \cong \varinjlim_n^{(\mathcal{C}/c)^{op}}(\Delta \mathbb{Z})$. So $H_q(t/c) = 0$ for $q > 0$ and $H_0(t/c) \cong \mathbb{Z}$. $\square$

Corollary 2.18 *Let $M(E, I)$ be a free partially commutative monoid which does not contain $> n$ mutually commuting generators. Then $\text{hd } M(E, I) \leq n$.*

PROOF. It follows from Lemma 2.17 for $\mathcal{C} = M(E, I)^{op}$ that the functor $t : \mathfrak{F}(M(E, I)^{op}) \rightarrow M(E, I)^{op}$ is strong coinital. The categories $\mathfrak{F}(M(E, I)^{op})$ and $\mathfrak{F}M(E, I)$ are isomorphic. Therefore $\text{hd}(\mathfrak{F}M(E, I))^{op} \geq \text{hd } M(E, I)$. By Corollary 2.10 we have $\text{hd } M(E, I) \leq n$. $\square$

Corollary 2.19 *Let $M(E, I)$ be a locally bounded free partially commutative monoid and G a right $M(E, I)$ -module. Then the homology groups $H_n(M(E, I)^{op}, G)$ are isomorphic to the homology groups of the chain complex*

$$\begin{aligned} 0 \leftarrow G \xleftarrow{d_1} \bigoplus_{a_1 \in E} G \xleftarrow{d_2} \bigoplus_{\substack{a_1 < a_2 \\ (a_1, a_2) \in I}} G \leftarrow \dots \\ \dots \leftarrow \bigoplus_{\substack{a_1 < a_2 < \dots < a_{n-1} \\ (a_i, a_j) \in I}} G \xleftarrow{d_n} \bigoplus_{\substack{a_1 < a_2 < \dots < a_n \\ (a_i, a_j) \in I}} G \leftarrow \dots, \end{aligned}$$

the n -th member of which for $n \geq 0$ is equal to the direct sum of copies of the group G by all sequences of mutually commuting generators $a_1 < a_2 < \dots < a_n$ belonging to E and the differential is defined by

$$d_n((a_1, \dots, a_n), g) = \sum_{s=1}^n (-1)^s ((a_1, \dots, \widehat{a_s}, \dots, a_n), ga_s - g)$$

where $ga_s \in G$ denotes a member obtained by right action of a_s on $g \in G$.

PROOF. By Lemma 2.17 $t : \mathfrak{F}M(E, I) \rightarrow M(E, I)$ is strong coinital. Thus the groups $H_n(M(E, I)^{op}, G)$, which equal $\varinjlim_n^{M(E, I)^{op}} G$ by definition, are isomorphic to $\varinjlim_n^{\mathfrak{F}M(E, I)^{op}} G \circ t^{op}$. Substituting $t(1, a_s) = a_s$, $t(a_s, 1) = 1$, $G \circ t^{op}(1, a_s)(g) = ga_s$, $G \circ t^{op}(a_s, 1)(g) = 1$, we obtain from Theorem 2.13 the required chain complex. $\square$

3 Homology of right $M(E, I)$ -sets with coefficients in diagrams

3.1 Homology groups of right $M(E, I)$ -sets

Theorem 3.1 *Let $M(E, I)$ be a locally bounded free partially commutative monoid, $X : M(E, I)^{op} \rightarrow \text{Set}$ a right $M(E, I)$ -set, $F : (M(E, I)/X)^{op} \rightarrow \text{Ab}$ a functor. Then the homology groups $\varinjlim_n^{(M(E, I)/X)^{op}} F$ are isomorphic to the homology groups of the chain complex*

$$\begin{aligned} 0 \leftarrow \bigoplus_{x \in X} F(x) \xleftarrow{d_1} \bigoplus_{(x, a_1)} F(x) \xleftarrow{d_2} \bigoplus_{\substack{(x, a_1 a_2), a_1 < a_2 \\ (a_1, a_2) \in I}} F(x) \leftarrow \dots \\ \dots \leftarrow \bigoplus_{\substack{(x, a_1 \dots a_{n-1}), a_1 < a_2 < \dots < a_{n-1} \\ (a_i, a_j) \in I}} F(x) \xleftarrow{d_n} \bigoplus_{\substack{(x, a_1 \dots a_n), a_1 < a_2 < \dots < a_n \\ (a_i, a_j) \in I}} F(x) \leftarrow \dots, \end{aligned}$$

with

$$d_n(x, a_1 \dots a_n, g) = \sum_{s=1}^n (-1)^s \left((x a_s, a_1 \dots \widehat{a_s} \dots a_n, F(x \xrightarrow{a_s} x \cdot a_s)g) - (x, a_1 \dots \widehat{a_s} \dots a_n, g) \right)$$

PROOF. In a similar manner with the proof of [12, Theorem 5.3] consider an arbitrary monoid M with functors $X : M^{op} \rightarrow \text{Set}$ and $F : (M/X)^{op} \rightarrow \text{Ab}$. Let $S : (M/X)^{op} \rightarrow M^{op}$ be a (dual of forgetful) functor which acts as $S(x \xrightarrow{\mu} y) = \mu$. For any functor $F : (M/X)^{op} \rightarrow \text{Ab}$ there is the left Kan extension $\text{Lan}^S F$. Each of the connected components of S/M where M denotes the unique object of the monoid considered as a small category has a terminal object of the form $(x, 1)$ for some $x \in X$. It follows from here that $\text{Lan}^S F$ is a right M -module $\bigoplus_{x \in X} F(x)$ with the action $(x, g) \cdot \mu = (x\mu, F(x \xrightarrow{\mu} x\mu)(g))$. Since the coproduct functor is exact in Ab , the functor Lan^S is exact. Moreover, Lan^S is left adjoint to the exact functor $(-) \circ S : \text{Ab}^{M^{op}} \rightarrow \text{Ab}^{(M/X)^{op}}$ and therefore it carries projective objects to projective. Consequently, Lan^S carries the projective resolution $0 \leftarrow F \leftarrow P_0 \leftarrow P_1 \leftarrow \dots$ of the object $F \in \text{Ab}^{(M/X)^{op}}$ to a projective resolution of $\text{Lan}^S F \in \text{Ab}^{M^{op}}$. Applying the functor $\varinjlim_n^{(M/X)^{op}}$ to the resolution of the object F we obtain a chain complex which homology groups are isomorphic

to $\varinjlim_n^{(M/X)^{op}} F$. It follows from an isomorphism $\varinjlim^{M^{op}} \text{Lan}^S \cong \varinjlim^{(M/X)^{op}}$ that this chain complex may be obtained by the application of the functor $\varinjlim^{M^{op}}$ to a projective resolution of the right M -module $\text{Lan}^S F$. Hence its the homology groups are isomorphic to $\varinjlim_n^{M^{op}} \text{Lan}^S F$. Consequently, the groups $\varinjlim_n^{(M/X)^{op}} F$ are isomorphic to the homology groups of M with coefficients in the right M -module $\text{Lan}^S F$.

Substitute $M = M(E, I)$. By Theorem 2.13 we obtain the chain complex for the calculating the homology groups with coefficients in $\text{Lan}^S F$

$$\begin{aligned} 0 \leftarrow \text{Lan}^S F \xleftarrow{d_1} \bigoplus_{a_1 \in E} \text{Lan}^S F \xleftarrow{d_2} \bigoplus_{\substack{a_1 < a_2 \\ (a_1, a_2) \in I}} \text{Lan}^S F \leftarrow \dots \\ \dots \leftarrow \bigoplus_{\substack{a_1 < a_2 < \dots < a_{n-1} \\ (a_i, a_j) \in I}} \text{Lan}^S F \xleftarrow{d_n} \bigoplus_{\substack{a_1 < a_2 < \dots < a_n \\ (a_i, a_j) \in I}} \text{Lan}^S F \leftarrow \dots, \end{aligned}$$

with differentials

$$d_n((a_1, \dots, a_n), g) = \sum_{s=1}^n (-1)^s ((a_1, \dots, \widehat{a_s}, \dots, a_n), g \cdot a_s - g).$$

We have $\text{Lan}^S F = \bigoplus_{x \in X} F(x)$ and the actions of d_n are defined by values at $(x, a_1 \dots a_n, g)$, $g \in F(x)$. Hence this chain complex is isomorphic to the chain complex which members and differentials are described in the theorem. $\square$

Examples of calculating the homology groups of $M(E, I)$ -sets.

EXAMPLE 3.1 Let $M(E, I)$ be a locally bounded free partially commutative monoid which acts on the one-point set $X = \{\star\}$. In this case by Theorem 3.1 the homology groups $\varinjlim_n^{(M(E, I)/X)^{op}} \Delta \mathbb{Z}$ are isomorphic to the homology of the chain complex

$$\begin{aligned} 0 \leftarrow \mathbb{Z} \xleftarrow{d_1} \bigoplus_{a_1} \mathbb{Z} \xleftarrow{d_2} \bigoplus_{\substack{a_1 < a_2 \\ (a_1, a_2) \in I}} \mathbb{Z} \leftarrow \dots \\ \dots \leftarrow \bigoplus_{\substack{a_1 < a_2 < \dots < a_{n-1} \\ (a_i, a_j) \in I}} \mathbb{Z} \xleftarrow{d_n} \bigoplus_{\substack{a_1 < a_2 < \dots < a_n \\ (a_i, a_j) \in I}} \mathbb{Z} \leftarrow \dots, \end{aligned}$$

where

$$d_n(a_1 \cdots a_n) = 0$$

Consequently, $\varinjlim_n^{(M(E,I)/\{\star\})^{op}} \Delta \mathbb{Z} \cong \mathbb{Z}^{p_n}$, where p_n is the number of subsets $\{a_1, \dots, a_n\} \subseteq E$ consisting of mutually commuting generators. (The number of empty subsets $p_0 = 1$.)

We will show that the homology groups of $M(E, I)$ -sets can have a torsion.

Recall that a *simplicial schema* is a pair $(X, \mathfrak{M})$ consisting of a set X with a set $\mathfrak{M}$ of its nonempty subsets for which the following implication holds

$$S \subseteq T \in \mathfrak{M} \Rightarrow S \in \mathfrak{M}.$$

Let $(X, \mathfrak{M})$ be a simplicial schema. Consider an arbitrary total order relation $<$ on X . Let

$$X_n = \{(x_0, x_1, \dots, x_n) : \{x_0, x_1, \dots, x_n\} \in \mathfrak{M} \text{ \& } x_0 < x_1 < \dots < x_n\}.$$

Consider the family of abelian groups

$$C_n(X, \mathfrak{M}) = \begin{cases} L(X_n), & \text{for } n \geq 0, \\ 0, & \text{otherwise.} \end{cases}$$

Define homomorphisms $d_n : C_n(X, \mathfrak{M}) \rightarrow C_{n-1}(X, \mathfrak{M})$ at elements of basis by

$$d_n(x_0, x_1, \dots, x_n) = \sum_{i=0}^n (-1)^i (x_0, \dots, x_{i-1}, x_{i+1}, \dots, x_n).$$

It is well known that $(C_n(X, \mathfrak{M}), d_n)$ is a chain complex. Denote $C_*(X, \mathfrak{M}) = (C_n(X, \mathfrak{M}), d_n)$. It easy to prove that homology groups of the chain complex $C_*(X, \mathfrak{M})$ do not depend on the order relation. These groups are called *homology groups* $H_n(X, \mathfrak{M})$ of the *simplicial schema* $(X, \mathfrak{M})$.

EXAMPLE 3.2 Let $M(E, I)$ be any locally bounded free partially commutative monoid with the right action on the set $X = \{x_0, \star\}$ by the laws

$$x_0 \cdot a = \star, \star \cdot a = \star \quad a \in E.$$

Let $\mathbb{Z}[x_0] : (M(E, I)/X)^{op} \rightarrow \text{Ab}$ be a functor such that $\mathbb{Z}[x_0](x_0) = \mathbb{Z}$ and $\mathbb{Z}[x_0](\star) = 0$. By Theorem 3.1 the groups $\varinjlim_n^{(M(E, I)/X)^{op}} \mathbb{Z}[x_0]$ are isomorphic to the homology groups of the chain complex

$$\begin{aligned} 0 \leftarrow \mathbb{Z} \xleftarrow{d_1} \bigoplus_{a_1} \mathbb{Z} \xleftarrow{d_2} \bigoplus_{\substack{a_1 < a_2 \\ (a_1, a_2) \in I}} \mathbb{Z} \leftarrow \dots \\ \dots \leftarrow \bigoplus_{\substack{a_1 < a_2 < \dots < a_{n-1} \\ (a_i, a_j) \in I}} \mathbb{Z} \xleftarrow{d_n} \bigoplus_{\substack{a_1 < a_2 < \dots < a_n \\ (a_i, a_j) \in I}} \mathbb{Z} \leftarrow \dots, \end{aligned}$$

where

$$d_n(a_1 \cdots a_n) = \sum_{s=1}^n (-1)^{s+1} (a_1 \cdots \widehat{a_s} \cdots a_n)$$

Consequently, $\varinjlim_n^{M(E, I)/X^{op}} \mathbb{Z}[x_0] \cong H_{n-1}(E, \mathfrak{M})$ for all $n \geq 2$.

Let $(M(E, I), X)$ be a pair consisting of a free commutative monoid and a right $M(E, I)$ -set X . Consider sets

$$Q_n X = \{(x, a_1 \cdots a_n) : a_1 < \cdots < a_n \& (1 \leq i < j \leq n \Rightarrow (a_i, a_j) \in I)\}$$

and maps

$$Q_n X \xrightleftharpoons[\partial_i^{n,1}]{\partial_i^{n,0}} Q_{n-1} X, \quad n \geq 1, \quad 1 \leq i \leq n, \quad \varepsilon \in \{0, 1\},$$

defined by

$$\partial_i^{n,0}(x, a_1 \cdots a_n) = (x, a_1 \cdots \widehat{a_i} \cdots a_n), \quad \partial_i^{n,1}(x, a_1 \cdots a_n) = (xa_i, a_1 \cdots \widehat{a_i} \cdots a_n).$$

Lemma 3.2 *The sets $Q_n X$ and the maps $\partial_i^{n,0}, \partial_i^{n,1}$ constitute a precubical set.*

PROOF. It may be directly checked that for every $x \in X$ and $\alpha, \beta \in \{0, 1\}$ there are the following equalities $\partial_i^{n-1,\alpha} \partial_j^{n,\beta}(x) = \partial_{j-1}^{n-1,\beta} \partial_i^{n,\alpha}(x)$, for $1 \leq i \leq n$. Hence $\partial_i^{n-1,\alpha} \partial_j^{n,\beta} = \partial_{j-1}^{n-1,\beta} \partial_i^{n,\alpha}$. $\square$

Corollary 3.3 *Let $X : M(E, I)^{op} \rightarrow \text{Set}$ be a right $M(E, I)$ -set. Then the homology groups $\varinjlim_n^{(M(E, I)/X)^{op}} \Delta \mathbb{Z}$ are isomorphic to integral homology groups of the precubical set Q_*X .*

PROOF. Substitute in Theorem 3.1 $F = \Delta \mathbb{Z}$. By Definition 1.1 the homology groups of the resulting chain complex $C_*(Q_*X)$ are precisely the homology groups of the precubical set Q_*X . $\square$

3.2 Homology of asynchronous transition systems

DEFINITION 3.3 [30] *An asynchronous transition system*

$$A = (S, s_0, E, I, \text{Tran})$$

is a data consisting of sets S and E , a member $s_0 \in S$, a subset $\text{Tran} \subseteq S \times E \times S$, and an irreflexive symmetric relation $I \subseteq E \times E$ satisfying to the following three conditions

- (i) *for each $e \in E$ there are $s, s' \in S$ such that $(s, e, s') \in \text{Tran}$;*
- (ii) *if $(s, e, s') \in \text{Tran}$ and $(s, e, s'') \in \text{Tran}$, then $s' = s''$;*
- (iii) *for every pair $(e_1, e_2) \in I$ and for arbitrary triples $(s, e_1, s_1) \in \text{Tran}$, $(s_1, e_2, u) \in \text{Tran}$ there exists $s_2 \in S$ such that $(s, e_2, s_2) \in \text{Tran}$ and $(s_2, e_1, u) \in \text{Tran}$.*

Members of S are called states, members of Tran are transitions, $s_0 \in S$ is an initial state, I is independence relation.

A morphism of asynchronous transition systems

$$(\eta, \sigma) : (S, s_0, E, I, \text{Tran}) \rightarrow (S', s'_0, E', I', \text{Tran}')$$

consists of a map $\sigma : S \rightarrow S'$ and a partial function $\eta : E \rightarrow E'$ for which $\sigma(s_0) = s'_0$ and the following two conditions hold

- (i) *$(s, a, t) \in \text{Tran} \Rightarrow (\sigma(s), \eta(a), \sigma(t)) \in \text{Tran}'$, if $\eta(a)$ is defined and otherwise $\sigma(s) = \sigma(t)$;*
- (ii) *if $(e_1, e_2) \in I$ and $\eta(e_1), \eta(e_2)$ are both defined, then $(\eta(e_1), \eta(e_2)) \in I'$.*

The category of asynchronous transition systems we denote by $\mathfrak{A}$.

Asynchronous transition systems and right pointed sets over monoids.

DEFINITION 3.4 A right pointed set over a monoid is a triple $(M, \cdot, X)$ consisting of a monoid M , a right pointed M -set X , and a map $\cdot : X \times M \rightarrow X$ such that

$$\begin{aligned} \star \cdot \mu &= \star \text{ for all } \mu \in M; \\ (x \cdot \mu_1) \cdot \mu_2 &= x \cdot (\mu_1 \mu_2) \text{ for all } x \in X, \mu_1, \mu_2 \in M; \\ x \cdot 1 &= x, \text{ for all } x \in X. \end{aligned}$$

Here 1 is the identity element of the monoid. The symbol $' \cdot '$ is omitted usually. So a right pointed set over a monoid is denoted by (M, X) .

A morphism of right pointed sets over monoids $(\eta, \sigma) : (M, X) \rightarrow (M', X')$ is a pair consisting of a monoid homomorphism $\eta : M \rightarrow M'$ and based map $\sigma : X \rightarrow X'$ satisfying $\sigma(x \cdot \mu) = \sigma(x) \cdot \eta(\mu)$ for all $x \in X$ and $\mu \in M$. Denote the category of right pointed sets over monoids by (Mon, Set_*) .

Define a map $Ob(\mathfrak{A}) \rightarrow (Mon, Set_*)$ assigning to every asynchronous transition system $(S, s_0, E, I, Tran)$ a pair $(M(E, I), S_*)$ where S_* is the pointed set with the action $(-) \cdot e : S \cup \{\star\} \rightarrow S \cup \{\star\}$ on generators $e \in E$ by

$$s \cdot e = \begin{cases} s' , & \text{if } (s, e, s') \in Tran, \\ \star , & \text{if there is not } s' \text{ such that } (s, e, s') \in Tran. \end{cases}$$

It is proved in [12] that the map $Ob(\mathfrak{A}) \rightarrow (Mon, Set_*)$ may be extended to a functor $\mathfrak{A} \rightarrow (Mon, Set_*)$. This functor assigns to any morphism

$$(\eta, \sigma) : (S, s_0, E, I, Tran) \rightarrow (S', s'_0, E', I', Tran')$$

of asynchronous transition systems the morphism $(\eta^*, \sigma_*) : (M(E, I), S_*) \rightarrow (M(E', I'), S'_*)$ consisting of an extension of the map $\eta : E \rightarrow E'$ to a monoid homomorphism and of natural transformation $\sigma_* : S_* \rightarrow S'_* \circ (\eta^*)^{op}$ consisting of the map

$$\sigma_*(s) = \begin{cases} \sigma(s) , & \text{if } s \in S, \\ \star , & \text{if } s = \star. \end{cases}$$

A forgetful functor $U : Set_* \rightarrow Set$ defined by $X \mapsto X$ and $f \mapsto f$. An augmented state category $K_*(T)$ [12, Definition 5.1] of an asynchronous transition system $T = (S, s_0, E, I, Tran)$ is the category which objects are the members $s \in S \cup \{\star\}$ and the morphisms are triples (μ, s, s') consisting of $\mu \in M(E, I)$, $s, s' \in S$ satisfying to $s \cdot \mu = s'$.

A homology groups $H_n(K_*(T), F)$ of an asynchronous transition system T with coefficients in a functor $F : K_*(T) \rightarrow Ab$ are abelian groups $\varinjlim_n^{K_*(T)} F$.

Events $e_j \in E$, $j \in J$ are said to be *mutually independent*, if $(e_j, e_{j'}) \in I$ for all $j, j' \in J$ such that $j \neq j'$.

Theorem 3.4 *Let $T = (S, s_0, E, I, Tran)$ be an asynchronous transition system without infinite sets of mutually independent events with an arbitrary total order relation $' < '$ on E and $F : K_*(T) \rightarrow \text{Ab}$ a functor. Then the homology groups $H_n(K_*(T), F)$ are isomorphic to homology groups of the following chain complex*

$$\begin{array}{ccccccc}
0 \leftarrow & \bigoplus_{s \in S \cup \{\star\}} & F(s) & \xleftarrow{d_1} & \bigoplus_{(s, e_1)} & F(s) & \xleftarrow{d_2} & \bigoplus_{\substack{(s, e_1 e_2), e_1 < e_2 \\ (e_1, e_2) \in I}} & F(s) & \leftarrow \dots \\
\dots & \leftarrow & \bigoplus_{\substack{(s, e_1 \dots e_{n-1}), e_1 < e_2 < \dots < e_{n-1} \\ (e_i, e_j) \in I}} & F(s) & \xleftarrow{d_n} & \bigoplus_{\substack{(s, e_1 \dots e_n), e_1 < e_2 < \dots < e_n \\ (e_i, e_j) \in I}} & F(s) & \leftarrow \dots
\end{array}$$

where

$$\begin{aligned}
d_n(s, e_1 \dots e_n, g) = \\
\sum_{i=1}^n (-1)^i \left((s \cdot e_i, e_1 \dots \widehat{e_i} \dots e_n, F(s \xrightarrow{e_i} s \cdot e_i)g) - (s, e_1 \dots \widehat{e_i} \dots e_n, g) \right)
\end{aligned}$$

PROOF. It easy to see that $K_*(T)$ is isomorphic to a category which is dual to $M(E, I)/U \circ S_*$. Therefore the wanted assertion follows from Theorem 3.1. $\square$

If the asynchronous transition system T has $\leq n$ mutually independent events, then the chain complex obtained in Theorem 3.4, has the length $\leq n$. Hence we get the following

Corollary 3.5 *Let $n > 0$ be the maximal number of mutually independent events in an asynchronous transition system T . Then $H_k(K_*(T), F) = 0$ for all $k > n$.*

This assertion gives the positive answer to my question [12, Open problem 2]. Theorem 3.4 leads to the algorithm of calculating the integral homology groups of an asynchronous transition system with finite sets S and E by the computations of the Smith normal form of the differentials d_n .

References

- [1] Baues H.-J., Wirsching G. *Cohomology of small categories.* // J. Pure Appl. Algebra. 1985. V.38, N 2/3. P.187–211.

- [2] Bednarczyk M. A. *Categories of Asynchronous Systems* Ph.D. Thesis, University of Sussex, report 1/88, 1988. 222 p. <http://www.ipipan.gda.pl/~marek>
- [3] Bucur I., Deleanu A. *Introduction to the theory of categories and functors*. London-New York-Sydney: Interscience, 1968.
- [4] Diekert V., Métivier Y. *Partial Commutation and Traces*. // Handbook of formal languages. V. 3. Springer-Verlag, 1997. P. 457–533.
- [5] Gabriel P., Zisman M. *Calculus of fractions and homotopy theory*. Berlin-Heidelberg-New York: Springer-Verlag, 1967.
- [6] Gaucher P. *Homotopy invariants of higher dimensional categories and concurrency in computer science*. // Math. Structures Comput. Sci. 2000. V.10, N 4. P. 481–524.
- [7] Gaucher P. *About the globular homology of higher dimensional automata*. // Cah. Topol. Geom. Differ. 2002. V.43, N2. P.107–156.
- [8] Goubault E. *The Geometry of Concurrency*. Ph.D. Thesis, Ecole Normale Supérieure, 1995. 349 p. <http://www.dmi.ens.fr/~goubault>
- [9] Havas G., Majewski B.S. *Integer Matrix Diagonalization*. // J. Symbolic Computation. 1997. V.24, N 3/4. P. 399–408.
- [10] Husainov A. *Homological dimension theory of small categories*. // J. Math. Sci., New York. 2002. V. 110, N 1. P. 2273–2321. <http://www.kluweronline.com>
- [11] Husainov A.A. *On the homology of monoids and distributed systems* // 5th International Algebraic Conference in Ukraine. Abstracts. Odessa, 2005.
- [12] Husainov A. *On the homology of small categories and asynchronous transition systems*. // Homology Homotopy Appl., 2004. V. 6, N 1. P. 439–471. <http://www.rmi.acnet.ge/hha>
- [13] Husainov A. A., Tkachenko V. V. *Homology groups of asynchronous transition systems*. Mathematical modeling and the near questions of mathematics. Collection of the scientific works. Khabarovsk: KhGPU,

2003. P. 23–33. (Russian)
http://www.knastu.ru/husainov_site/index.html
- [14] Kaczynski T., Mischaikov K., Mrozek M. *Algebraic Topology: A Computational Approach*. Jagellonian University, 2000.
 - [15] Kaczynski T., Mischaikov K., Mrozek M. *Computational homology*. // Homology Homotopy Appl. 2003. V.5, N 2. P. 233 – 256.
 - [16] Khusainov A. A. *Cohomology of small categories with coefficients in an Abelian category with exact products*. // Sib. Math. J. 1989. V.30, N 4. P.668–672; translation from Sib. Mat. Zh. 1989. V. 30, N 4. P. 210–215.
 - [17] Khusainov A. A. *Homotopical equivalence of coverings and the spectral sequence of a fibration* // Sib. Math. J. 1991. V.32, N 1. P. 116-122; translation from Sib. Mat. Zh. 1991. V.32, N 1 (185). P.141-147.
 - [18] Khusainov A. A. *Comparison of the dimensions of a small category*. // Sib. Math. J. 1997. V. 38, N 6. P. 1230–1240. translation from Sib. Mat. Zh. 1997. V. 38, N 6. P. 1413–1426.
 - [19] Leech J. *Cohomology theory for monoid congruences*. // Houston J. Math. 1985. V.11, N 2. P. 207 – 223.
 - [20] MacLane S. *Categories for the Working Mathematician*. New York-Heidelberg-Berlin: Springer-Verlag, 1972 (Graduate Texts in Mathematics 5).
 - [21] Mac Lane S. *Homology*. Berlin-Göttingen-Heidelberg: Springer-Verlag, 1963.
 - [22] Mitchell B. *A Remark on Projectives in Functor Categories*. // J. Algebra. 1981. V.69, N 1. P.24-31.
 - [23] Mitchell B. *Rings with several objects*. // Adv. Math. 1972. V.8. P.1-161.
 - [24] Naudin P., Quitté C. *Algorithmique algébrique. Avec exercices corrigés*. Paris-Milan-Barcelone-Bonn: Masson, 1992.
 - [25] Oberst U. *Basisweiterung in der Homologie kleiner Kategorien* // Math.Z. 1967. Bd.100. S.36–58.

- [26] Oberst U. *Homology of categories and exactness of direct limits.* // Math. Z. 1968. Bd. 107. S.87–115.
- [27] Polyakova L.Yu. *Resolutions for free partially commutative monoids* // Preprint, 2006. 10p. <http://arxiv.org/abs/math.KT/0606746>
- [28] Schubert H. *Kategorien II.* Berlin: Akademie-Verlag, 1970.
- [29] Seifert H., Threlfall W. *Lehrbuch der Topologie.* Leipzig: Teubner, 1934.
- [30] Winskel G., Nielsen M. *Models for Concurrency.* // Handbook of Logic in Computer Science, Vol. IV, ed. Abramsky, Gabbay and Maibaum. Oxford University Press, 1995. P.1–148.